

Study finds simple interventions reduce newborn deaths in Africa

February 3 2011

Training community birth attendants in rural Zambia in a simple newborn resuscitation protocol reduced neonatal deaths by nearly 50 percent - a finding that shows high potential to save lives in similar remote settings, a team of Boston University School of Public Health [BUSPH] international health researchers is reporting.

Findings published Feb. 3 in the *BMJ* by the team from the BU Center for Global Health and Development show that training and equipping Zambian traditional birth attendants to perform a neonatal resuscitation intervention led to a net reduction of about 18 deaths per 1,000 live births in the first 28 days of life - a significant reduction in the overall neonatal mortality rate in [Zambia](#), of about 34 deaths per 1,000 live births.

"We essentially showed that there is a method, very simple and inexpensive, to reduce neonatal mortality by about half," said Dr. Christopher Gill, associate professor of international health at BUSPH and lead author on the study. "This is the first high-quality study, using a randomized control methodology, to show unambiguously that a neonatal resuscitation protocol can be used to save lives. It's hugely encouraging."

The study, dubbed "LUNESP" for the Lufwanyama Neonatal Survival Project, was a large-scale trial in the rural Lufwanyama district of Zambia, where access to healthcare facilities is extremely limited. Researchers worked with Zambian traditional birth attendants - community members who attend to villagers' births - to gauge the impact

of a package of neonatal interventions.

Birth attendants were divided into two groups - one which used the existing standard-of-care in attending to deliveries, and the other that received extra training aimed at preventing neonatal hypothermia and birth asphyxia. Globally, at least 10 percent of otherwise healthy newborns experience inadequate respiration at birth, a problem often compounded by hypothermia from failure to immediately dry and warm the infants.

The trained attendants were instructed to immediately dry and swaddle the newborn in a dry blanket; to suction the infant's mouth and nose with a rubber suction bulb; and, when indicated, to stimulate breathing by rubbing the infant's feet or back or, in cases of inadequate respiration, to use a tube-mask to resuscitate the infant.

Non-trained attendants followed traditional practices, wrapping the infant in thin cloth and paying minimal attention to clearing the airways, sometimes by wiping the infant's nose with a cloth.

Among 3,497 deliveries, mortality in the first 24 hours of life was significantly lower for those delivered by trained attendants - 7.8 deaths per 1,000 live births, compared to 19.9 per 1,000 births for the non-trained group. Deaths due to birth asphyxia were reduced by 63 percent among infants delivered by the trained attendants, the study found.

"Given recent estimates that approximately 800,000 infants die each year due to birth asphyxia, these findings have broad public health relevance," the research team said.

A second intervention involved training the attendants to recognize symptoms of sepsis in the first four weeks of life, and to administer a dose of oral amoxicillin to infants before referring them to a health

center for further care. While this intervention was found to be effective -- with lower rates of neonatal mortality during weeks 2-4 of an infant's life -- the difference did not reach statistical significance.

"While early identification and treatment of sepsis is important, we found that most of the deaths were prevented in the first hours of life," Gill said.

The study provided evidence that community members who serve as birth attendants can perform a vital role in access to health care.

"In the context of a highly dispersed, rural African community with very limited access to health care, traditional birth attendants [TBAs] were able to master a set of skills that allowed them to significantly reduce neonatal mortality," the study concludes. "This was accomplished in a population of women TBAs with very low rates of formal education and under austere conditions. . . We believe this approach has high potential to be applied in other resource-constrained settings."

Since the study ended, the organization Save the Children has stepped in with financial support to help to continue the interventions in Zambia, Gill said.

The LUNESP study comes on the heels of a recent published study [known as "First Breath"] in the New England Journal of Medicine that found little impact of a neonatal resuscitation protocol on mortality rates. The BUSPH researchers noted several differences between that study and their own, including that a third of deliveries in the First Breath study occurred in clinics or hospitals, and that the gap in training levels between birth attendants in that study was not as pronounced as in LUNESP.

"LUNESP and 'First Breath' addressed fundamentally different

questions, in different populations, using very different methods," the BUSPH team wrote.

The BU Center for Global Health and Development is involved in a variety of research studies aimed at improving infant and child survival by training and empowering community health workers in Africa and other regions of the world.

Provided by Boston University Medical Center

Citation: Study finds simple interventions reduce newborn deaths in Africa (2011, February 3) retrieved 20 July 2023 from <https://medicalxpress.com/news/2011-02-simple-interventions-newborn-deaths-africa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.