

Social norms for obesity learned in childhood

December 19 2011

Newcastle University research studying siblings has revealed that childhood experience and genes may set your weight rather than social networks later in life.

An individual's social norms in terms of what they consider acceptable weight are formed during childhood, and are not highly influenced by our later social networks according to the study published online this week in <u>Obesity</u>.

Dr Heather Brown and colleagues investigated the effects of timeconstant factors such as genetics and upbringing, and changeable factors, like friends and opportunities for exercise, on the development of social norms regarding weight in children.

They used data from the Panel Study of Income Dynamics to analyze the correlation in sibling body mass index (BMI) data from 236 adolescent siblings living together, and 838 adult siblings living apart.

Time-constant factors significantly influenced both groups, but only the adolescent group was influenced by changeable factors. A larger study may provide findings that are more robust.

The authors conclude that previous studies may have overestimated the effect of social networks on BMI and Dr Brown and colleagues suggest that if obesity prevention strategies are to be effective, they should be implemented in early childhood and at a household level.



More information: Exploring the Factors Contributing to Sibling Correlations in BMI: A Study Using the Panel Study of Income Dynamics, <u>DOI:10.1038/OBY.2011.351</u>

Provided by Newcastle University

Citation: Social norms for obesity learned in childhood (2011, December 19) retrieved 24 March 2023 from <u>https://medicalxpress.com/news/2011-12-social-norms-obesity-childhood.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.