

Secondhand smoke exposure linked to signs of heart disease

March 7 2013

Nonsmokers, beware. It seems the more you are exposed to secondhand tobacco smoke—whether it was during your childhood or as an adult, at work or at home—the more likely you are to develop early signs of heart disease, according to research being presented at the American College of Cardiology's 62nd Annual Scientific Session.

Researchers found that 26 percent of people exposed to varying levels of [secondhand smoke](#) had signs of coronary artery calcification (CAC), compared to 18.5 percent in the general population. The new data also shows that people who report higher levels of secondhand [tobacco smoke exposure](#) also have the greatest evidence of coronary artery calcification, a build-up of calcium in the [artery walls](#) as seen on a low-dose computed tomography scan. While previous studies have shown a marked increase in [cardiac events](#) related to secondhand [tobacco smoke](#), authors say this study is the first to demonstrate a clear dose-response relationship between [secondhand smoke exposure](#) and the earliest detectable signs of heart disease.

"This research provides additional evidence that secondhand smoke is harmful and may be even more dangerous than we previously thought," said Harvey Hecht, MD, associate director of [cardiac imaging](#) and professor of medicine at Mount Sinai Medical Center and study author. "We actually found the risk of secondhand [smoke exposure](#) to be an equivalent or stronger risk factor [for CAC] than other well-established ones such as [high cholesterol](#), hypertension and diabetes. Passive exposure to smoke seems to independently predict both the likelihood

and extent of CAC."

After adjusting for other [cardiovascular risk factors](#), people classified as having low, moderate or high secondhand smoke exposure were 50, 60 and 90 percent more likely to have evidence of [coronary artery calcification](#) than those who reported minimal exposure. The apparent health effects of secondhand smoke on CAC remained regardless of whether the exposure was during childhood or adulthood. Dr. Hecht said these results further underscore the need for enforceable public smoking bans and other measures to reduce passive inhalation of cigarette smoke.

"Tobacco smoke can damage the coronary arteries of nonsmokers through many different ways, which can lead to plaque formation and then to heart attacks, so this lends more [credence] to enforcing smoking bans," Dr. Hecht said. "We know heart disease is significantly accelerated by secondhand smoke exposure, so it should be included as a routine part of medical exams and discussions about heart disease and try to prevent it as best we can."

This study included 3,098 healthy people between 40 and 80 years old who had never smoked (defined as having smoked fewer than 100 cigarettes in their lifetime) and who were already enrolled in the Flight Attendant Medical Research Institute (FAMRI)-International Early Lung Cancer Action Program CT screening program from 2005-2012. Each provided detailed health information and had a single low-dose non-gated CT scan as part of the FAMRI program. A questionnaire was administered to gauge personal exposures to secondhand smoke. Researchers then assigned a total secondhand tobacco smoke exposure score based on a how long they were exposed (years), daily intensity (packs per day) and if smokers were allowed to smoke anywhere or if they were restricted or not permitted (permission status).

Compared to those who had minimal exposure to secondhand smoke,

people with higher levels of exposure tended to be older, were more often women and more likely to have diabetes, high cholesterol and high blood pressure.

Although his team did not use the standard Agatston score to assess CAC, Dr. Hecht said this study further validates the utility of low-dose non-gated CT scans to measure the amount of plaque in the coronary arteries in nonsmokers exposed to secondhand tobacco smoke.

"By using this [imaging/approach], people who have been exposed to tobacco smoke can be evaluated for lung cancer, emphysema and coronary artery disease in a single low-dose scan," he said. As a general rule, people should limit secondhand smoke exposure as much as possible.

Future research directions will include how to determine if decreasing exposure to secondhand tobacco smoke can reduce new plaque formation and/or cardiac events in this group. This study was funded by the Flight Attendants Medical Research Institute.

More information: This study will be published online in the *Journal of the American College of Cardiology: Cardiovascular Imaging*.

Dr. Hecht will present the study "Secondhand Tobacco Smoke in Never Smokers is a Significant Risk Factor for Coronary Artery Calcification" on Sunday, March 10 at 11:45 a.m., Moscone Center, West, Room 3004.

Provided by American College of Cardiology

Citation: Secondhand smoke exposure linked to signs of heart disease (2013, March 7) retrieved 31 January 2024 from <https://medicalxpress.com/news/2013-03-secondhand-exposure-linked->

[heart-disease.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.