

Pilot study finds ER patients drinking high-octane beer

August 14 2013

Budweiser, Steel Reserve, Colt 45, Bud Ice and Bud Light – were consumed in the highest quantities by emergency room patients, according to a new pilot study from researchers at The Center on Alcohol Marketing and Youth (CAMY) at the Johns Hopkins Bloomberg School of Public Health. Three of these are "malt liquors" with higher alcohol content than regular beer.

The pilot study, published by *Substance Use and Misuse*, is the first study of its kind to assess [alcohol consumption](#) by brand and type from patients reporting to the [emergency department](#) with injury.

"Recent studies reveal that nearly a third of injury visits to Level I [trauma centers](#) were [alcohol](#)-related and frequently a result of heavy drinking," said lead study author David Jernigan, PhD, CAMY director. "Understanding the relationship between alcohol brands and their connection to injury may help guide policy makers in considering taxation and physical availability of different types of alcohol given the harms associated with them."

The study was conducted in an urban medical center at the Johns Hopkins Hospital Emergency Department in East Baltimore on Friday and Saturday nights between April 2010 and June 2011. Of the 105 respondents who admitted to drinking alcohol before their injury, 73 (69%) were male, and 72 (69%) were African American, reflecting the demographic profile of the neighborhood in which the emergency department is located.

The research team also tracked the ER patients' consumption of alcohol by type and compared it to national market share data from Impact Databank, a market research firm that tracks the U.S. market for alcoholic beverages by type and brand. The study found that the proportion of distilled spirits consumed by the ER sample was higher than the market share for distilled spirits in the U.S. More specifically, vodka, gin and brandy/cognac were over-represented compared to their market share in the national distilled spirits market. The same was true for 'ready-to-drink' beverages (RTDs). Women in the ER sample were more likely to report consuming higher quantities of RTDs.

Although [beer](#) was consumed at a lower proportion in the ER sample compared to the proportion of its consumption in the national [market share](#) for beer, men in the ER sample were more likely to report consuming higher quantities of beer or malt liquors, which has higher [alcohol content](#) than regular beer.

Four malt liquors - Steel Reserve, Colt 45, Bud Ice and King Cobra - accounted for almost 50 percent (46%) of the beer consumed by the sample. Yet these four beverages accounted for only 2.4 percent of beer consumption in the general population.

The next step, according to study authors, would be to pursue this type of research be further explored in a larger sample of emergency department admissions for injury, across multiple cities and hospitals. Policy implications of this kind of research could include requirements for clear labeling of alcohol content on malt beverage containers, including serving size labeling; limits on malt liquor availability and marketing; and graduated taxation of beer based on alcohol content to discourage consumption of higher-alcohol products.

Provided by Johns Hopkins University Bloomberg School of Public

Health

Citation: Pilot study finds ER patients drinking high-octane beer (2013, August 14) retrieved 20 May 2023 from <https://medicalxpress.com/news/2013-08-er-patients-high-octane-beer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.