

Researchers discover that the body clock may influence morning peak in adverse cardiovascular events

November 11 2013

Cardiovascular disease is the number one cause of death in both men and women, and most adverse cardiovascular events tend to happen in the morning. In new findings published in the November issue of *Blood*, researchers from Brigham and Women's Hospital (BWH) and Oregon Health & Science University have discovered that the internal body clock may contribute to the morning peak in heart attacks and ischemic strokes.

"Our findings suggest that the circadian system, or the internal body clock, contributes to the increased risk for [cardiovascular events](#) in the morning," said Frank A.J.L. Scheer, PhD, director of the Medical Chronobiology Program at BWH and corresponding author of this study.

The researchers studied 12 healthy adult volunteers in the intensive physiological monitoring laboratories at BWH. Participants were assessed throughout a two-week laboratory protocol designed to desynchronize daily behavioral and environmental rhythms from internal [circadian rhythms](#).

Researchers specifically evaluated the role of Plasminogen activator inhibitor-1 (PAI-1), which inhibits the breakdown of [blood](#) clots and is thus a risk factor for blood clotting, one of the major contributors to heart attack and ischemic stroke. The researchers sought out to test whether this morning peak in PAI-1 is caused by the internal circadian

system or by behaviors that typically occur in the morning, such as altered posture and physical activity. The researchers found a robust circadian rhythm in circulating PAI-1 with a peak corresponding to approximately 6:30 a.m. in a regular sleep/wake cycle.

"Our findings indicate that the human circadian system causes a morning peak in circulating levels of PAI-1, independent of any behavioral or environmental influences," explained Steven Shea, PhD, director of the Oregon Institute of Occupational Health Sciences and the co-author of this paper. "Indeed, the [circadian system](#) determined to a large extent the PAI-1 rhythm observed during a regular sleep/wake cycle. This morning peak in PAI-1 could help explain adverse cardiovascular events in vulnerable individuals."

The researchers added that these studies established the circadian control of PAI-1 in healthy individuals and that future research is required to test whether this rhythm is amplified, blunted or shifted in vulnerable individuals, such as with obesity, diabetes, or [cardiovascular disease](#).

Provided by Brigham and Women's Hospital

Citation: Researchers discover that the body clock may influence morning peak in adverse cardiovascular events (2013, November 11) retrieved 29 January 2023 from <https://medicalxpress.com/news/2013-11-body-clock-morning-peak-adverse.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--