

Postmenopausal breast cancer risk decreases rapidly after starting reg. physical activity

August 11 2014

Postmenopausal women who in the past four years had undertaken regular physical activity equivalent to at least four hours of walking per week had a lower risk for invasive breast cancer compared with women who exercised less during those four years, according to data published in *Cancer Epidemiology, Biomarkers & Prevention*, a journal of the American Association for Cancer Research.

"Twelve MET-h [metabolic equivalent task-hours] per week corresponds to walking four hours per week or cycling or engaging in other sports two hours per week and it is consistent with the World Cancer Research Fund recommendations of walking at least 30 minutes daily," said Agnès Fournier, PhD, a researcher in the Centre for Research in Epidemiology and Population Health at the Institut Gustave Roussy in Villejuif, France. "So, our study shows that it is not necessary to engage in vigorous or very frequent activities; even [walking](#) 30 minutes per day is beneficial."

Postmenopausal women who in the previous four years had undertaken 12 or more MET-h of physical activity each week had a 10 percent decreased risk of invasive breast cancer compared with women who were less active. Women who undertook this level of physical activity between five and nine years earlier but were less active in the four years prior to the final [data](#) collection did not have a decreased risk for invasive breast cancer.

"Physical activity is thought to decrease a woman's risk for breast cancer

after menopause," said Fournier. "However, it was not clear how rapidly this association is observed after regular physical activity is begun or for how long it lasts after regular exercise stops.

"Our study answers these questions," Fournier continued. "We found that recreational physical activity, even of modest intensity, seemed to have a rapid impact on breast cancer risk. However, the decreased breast cancer risk we found associated with physical activity was attenuated when activity stopped. As a result, postmenopausal women who exercise should be encouraged to continue and those who do not exercise should consider starting because their risk of breast cancer may decrease rapidly."

Fournier and colleagues analyzed data obtained from biennial questionnaires completed by 59,308 [postmenopausal women](#) who were enrolled in E3N, the French component of the European Prospective Investigation Into Cancer and Nutrition (EPIC) study. The mean duration of follow-up was 8.5 years, during which time, 2,155 of the women were diagnosed with a first primary [invasive breast cancer](#).

The total amount of self-reported recreational physical activity was calculated in MET-h per week. The [breast cancer](#) risk-reducing effects of 12 or more MET-h per week of recreational [physical activity](#) were independent of body mass index, weight gain, waist circumference, and the level of activity from five to nine years earlier.

Provided by American Association for Cancer Research

Citation: Postmenopausal breast cancer risk decreases rapidly after starting reg. physical activity (2014, August 11) retrieved 4 July 2023 from <https://medicalxpress.com/news/2014-08-postmenopausal-breast-cancer-decreases-rapidly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.