

# Opioid pain reliever abuse called a top 5 public health challenge

February 24 2015, by Stasia Thompson

---

The Centers for Disease Control calls prescription painkiller abuse "one of the worst drug overdose epidemics in history."

New studies on [prescription painkillers](#) show that from 1999 to 2011, the consumption of hydrocodone more than doubled and oxycodone use increased by 500 percent. During that time, opioid pain reliever (OPR) overdose nearly quadrupled.

"The rise in opioid consumption has resulted in a doubling in visits to the emergency department for nonmedical OPR use but I also see patients who make errors with medications they are legitimately supposed to be taking," says Megan Rech, [emergency medicine](#) pharmacist, Loyola University Health System. Dr. Rech is a registered and licensed pharmacist with advanced residency training in critical care and is a dedicated pharmacist for Loyola's [emergency department](#).

Addicts will frequent emergency rooms complaining of phantom conditions to try to get painkillers. "At Loyola, we limit painkiller prescriptions to last between 7 and 10 days with no refills, to avoid abuse and to make sure people who are genuinely ill see their doctor for follow-up care," says Dr. Rech. "At Loyola, we utilize an Illinois database that tracks scheduled prescriptions to help identify OPR addicts."

Addiction is defined as continued use of a drug despite negative consequences.

"Red flags that a patient may be an addict include complaining of general pain or excess pain compared to the exam, refusing to see a specialist or to contact his/her primary care physician, asking for brand name painkillers, or displaying an overfamiliarity with opioids," says Dr. Rech.

Loyola emergency medicine physicians often will opt for non-opioid forms of pain relief. As an Emergency Department pharmacist, Dr. Rech screens appropriate patients for presence of any medication or substances and counsels staff and patients on prescriptions.

Dr. Rech is also careful to stress the importance of patient education. "Many patients are also unaware that they can build up a tolerance to medication. If they stop taking prescribed painkillers for a while, the tolerance decreases," says Dr. Rech. "A patient can then restart the medication at the previous dose, which is now too high, increasing the risk of an overdose."

"When physicians prescribe medication, they need to be very clear with the patient on what it is, when it should be taken, when it should be stopped and disposed of and if it is habit-forming," says Dr. Rech. "Some patients make honest mistakes with self-dosage that unfortunately result in overdose."

Provided by Loyola University Health System

Citation: Opioid pain reliever abuse called a top 5 public health challenge (2015, February 24) retrieved 30 January 2024 from <https://medicalxpress.com/news/2015-02-opioid-pain-reliever-abuse-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.