

# Ultrasonic surgery reduces pain and swelling after chin surgery

March 11 2016

---

For patients undergoing plastic surgery of the chin (genioplasty), the use of ultrasonic "piezosurgery" equipment reduces trauma, pain, and swelling, compared to traditional surgical drills, reports a study in the *The Journal of Craniofacial Surgery*.

"Piezosurgery may be a viable alternative to traditional osteotomy [bone-cutting] technique, as it reduces the degree of inflammation, pain, swelling, and morbidity, improving satisfaction and patient comfort," according to the report by Dr. Gilberto Sammartino of University of Naples Federico II, Italy, and colleagues.

## **Piezosurgery versus Traditional Instruments for Genioplasty**

The researchers compared complications after genioplasty performed using piezosurgery devices or traditional rotating drills. Genioplasty, sometimes called mentoplasty, is a [plastic surgery](#) procedure done to reshape the chin—for example, augmenting or reducing it—for cosmetic and/or functional reasons.

Piezosurgery is a relatively new approach that uses ultrasonic energy, rather than conventional surgical instruments, for cutting of bone.

"Several studies have demonstrated that bone healing using piezosurgery is more rapid than other techniques using drills or burs, thanks to a lower inflammatory bone response," Dr. Sammartino and coauthors write.

The study included 40 patients scheduled for genioplasty, as a primary procedure or after corrective jaw [surgery](#). Patients were randomly assigned to undergo genioplasty using either ultrasonic piezosurgery instruments or traditional drills. Pain, healing, and complications were compared from one to 15 days after surgery.

The results showed lower pain scores for patients undergoing piezosurgery, although the difference was significant only on the third and seventh day after surgery. Swelling also seemed to be reduced with piezosurgery, compared to cutting drills.

Both groups had reduced feeling in the chin area throughout the first 15 days after surgery, mainly due to nerve stretching. By six months, sensation normalized within six months for all [patients](#) in both groups. Pain and swelling were completely resolved as well.

Previous studies have shown that piezosurgery leads to better control of the inflammatory bone response induced by surgery, and less cell damage leading to increased [bone](#) remodeling after surgery. Dr. Sammartino and colleagues conclude, "Bone undergoes less stress during surgery and thus less pain and [swelling](#) postoperatively, which is in agreement with the results found in our trial: [pain](#) and discomfort were minimal compared to the traditional technique (saw and drills) especially in the immediate postoperative period of healing (within 3 days)."

**More information:** Rosario Rullo et al. The Use of Piezosurgery in Genioplasty, *Journal of Craniofacial Surgery* (2016). [DOI: 10.1097/SCS.0000000000002473](#)

Provided by Wolters Kluwer Health

Citation: Ultrasonic surgery reduces pain and swelling after chin surgery (2016, March 11)  
retrieved 23 February 2023 from <https://medicalxpress.com/news/2016-03-ultrasonic-surgery-pain-chin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.