

# Study suggests expanded range for emerging tick-borne disease

February 16 2018, by Elisabeth Reitman

---



Human cases of *Borrelia miyamotoi*. Credit: Michael Greenwood

Human cases of *Borrelia miyamotoi*, a tick-borne infection with some similarities to Lyme disease, were discovered in the eastern United States less than a decade ago. Now new research led by the Yale School of Public Health strongly suggests that the pathogen's geographic range is bicoastal, far larger than previously believed.

A scientific team headed by Peter Krause, M.D., and including Erol Fikrig, M.D., and Sukanya Narasimhan, Ph.D., both of the Yale School of Medicine, and Robert Lane, Ph.D. of the University of California, Berkeley, tested archived sera and found evidence that the [disease](#) exists in a Lyme disease endemic community in northern California. A previous study by Krause and colleagues published last year found evidence that the disease exists in Manitoba, Canada.

"*B. miyamotoi* infection causes a viral like illness and can cause relapsing fever. Health professionals in areas where *B. miyamotoi* infections are endemic need to be aware of its presence," said Krause, a senior research scientist at the Yale School of Public Health. "It is likely that we will discover that its range extends into many other regions of the United States, possibly wherever Lyme disease is found."

The study, published in the journal *PLOS One*, analyzed sera from 101 residents in each of two consecutive years. The sera of 2 residents in the first year and 7 residents in the second year tested positive for the presence of *B. miyamotoi* antibodies.

Krause noted that while the results strongly implicate the pathogen's presence, the laboratory results are not definitive. The test used also detects the presence of two other relapsing fever diseases related to *B. miyamotoi*. Thus, the exact cause of infection in the study subjects could not be determined conclusively, although epidemiological and clinical data strongly pointed to *B. miyamotoi* infection. Krause said these findings also illustrate the need to develop improved serodiagnostic assays capable of differentiating all *Borrelia* tick-borne infections.

*B. miyamotoi* was first identified in ticks in 1994 in Japan and then in Connecticut in 2000 by Durland Fish, Ph.D., at the Yale School of Public Health. Human cases were first reported in 2011 in Russia by Russian scientists and Durland Fish, now a professor emeritus at the

Yale School of Public Health, and Krause. Human *B. miyamotoi* infection subsequently was reported in the eastern United States, and parts of Europe and Japan. Many symptoms of the disease are similar to those of Lyme disease, but patients with *B. miyamotoi* infection also may experience relapsing fever. People with a compromised immune system, such as cancer patients who are receiving chemotherapy, may develop *B. miyamotoi* meningoencephalitis. Antibiotic treatment for the newly discovered infection is the same as for Lyme disease and is highly effective, Krause said.

In 2014 Krause, Fikrig and Narasimhan, and other colleagues found that about 4 percent of 639 healthy people living in southern New England had evidence of previous *B. miyamotoi* infection, compared to about 10 percent with evidence of previous Lyme disease infection.

"Although not as common as Lyme disease, the frequency of *B. miyamotoi* [infection](#) in southern New England appears to be comparable to that of other infections transmitted by the black-legged tick, such as anaplasmosis and babesiosis," said Krause.

**More information:** Peter J. Krause et al. Human *Borrelia miyamotoi* infection in California: Serodiagnosis is complicated by multiple endemic *Borrelia* species, *PLOS ONE* (2018). [DOI: 10.1371/journal.pone.0191725](#)

Provided by Yale University

Citation: Study suggests expanded range for emerging tick-borne disease (2018, February 16) retrieved 4 July 2023 from <https://medicalxpress.com/news/2018-02-range-emerging-tick-borne-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.