


Abiraterone could halve risk of prostate cancer death for some patients

December 28 2021



Credit: CC0 Public Domain

Adding abiraterone to the standard treatment for locally advanced prostate cancer, where the cancer has a high chance of spreading, could halve the risk of death from the disease.

The researchers suggest that hormone therapy using abiraterone with prednisolone, could significantly reduce [prostate cancer](#) deaths and

improve outcomes for thousands of people every year.

Using abiraterone for this group of people is now being considered for use in NHS England based on this research. If successful, it could be rolled out to patients immediately.

The study, published today in *The Lancet*, is part of the STAMPEDE trial and was led by a team at University College London and The Institute of Cancer Research, London and funded by Cancer Research UK and the MRC. The study followed patients over a six-year period.

Abiraterone for earlier-stage prostate cancer

More than 52,000 people are diagnosed with [prostate](#) cancer every year in the UK, and around 22,000 patients are considered at "high-risk" of their cancer spreading.

Abiraterone is currently being used for patients with advanced prostate cancer which has spread to other parts of the body. It's also given to men who have stopped responding to standard hormone treatment. The drug is given in combination with a steroid called prednisolone.

However, there have remained questions over its benefits and impact on survival in earlier-stage disease.

A total of 1974 patients were enrolled across two arms of the trial. 988 were given the current standard treatment, while 986 patients were given the [standard treatment](#) combined with abiraterone. Around half of those in the abiraterone group were also given enzalutamide, another hormone therapy.

'Could extend lives and prevent cancer from spreading'

After 6 years of monitoring, adding abiraterone alone, or with enzalutamide to standard prostate cancer treatment improved survival and decreased the chance of the cancer spreading. 7 percent of people receiving abiraterone died from prostate cancer during the 6-year follow up period, compared with 15 percent of those receiving standard care.

Around half of the people in the abiraterone group (527/986 men) also received enzalutamide, another type of hormone drug. However, using this combination of drugs did not further improve outcomes beyond those receiving abiraterone on its own, and caused an increase in side effects.

These results indicate that using abiraterone to treat earlier stage prostate cancer could extend lives and prevent the cancer from spreading.

The STAMPEDE trial is based at the MRC Clinical Trials Unit at UCL and led in collaboration with researchers at UCL Cancer Institute and The Institute of Cancer Research, London.

Researchers at The Institute of Cancer Research (ICR) with funding from Cancer Research UK discovered abiraterone and developed it with colleagues at The Royal Marsden. In 2012, Cancer Research UK lobbied to make abiraterone available on the NHS and its now used to treat thousands of patients with prostate cancer that has spread.

Preventing unnecessary deaths

Study co-leader Professor Nick James, Professor of Prostate and Bladder Cancer Research at The Institute of Cancer Research, London, and chief investigator of the STAMPEDE trial, said:

"STAMPEDE continues to deliver practice-changing results. Currently, abiraterone is only given to patients with very [advanced prostate cancer](#).

Our latest findings are the first to show the drug can also benefit men whose cancer is at an earlier stage—improving survival and reducing the chance of progression. The next step is for NICE to consider and implement our findings, so that men can benefit from [abiraterone](#) before their cancer has spread, drastically improving their quality of life and preventing many unnecessary deaths."

Study co-leader Professor Gert Attard, UCL Cancer Institute said:

"This is the first time we've seen a treatment for this kind of prostate cancer that can do more than extend life. We're seeing clear and convincing evidence that some people who would have died of prostate cancer, the third leading cause of cancer death in the UK, will no longer die from it."

Yet more people could soon benefit from this research

Michelle Mitchell, Chief Executive of Cancer Research UK, said:

"These results are the latest in a long line of practice changing findings from our STAMPEDE trial. It's recruited over 10,000 patients and has led to 29 changes in clinical practice across the world, directly influencing the treatment of people with prostate cancer. It's great to see that yet more people with prostate cancer could soon see benefit from this innovative research."

Ken was diagnosed with prostate cancer in 2015 and was offered the chance to be part of a separate part of the STAMPEDE trial, looking at a hormone treatment for prostate cancer called Zoladex. He said:

"My treatment would not have happened but for someone before me taking part in a clinical trial, so I just wanted to give something back. Knowing that people with prostate [cancer](#) might have better odds in the

future in part because of what they learnt from my trial feels like I've paid it forward."

Five years down the line, Ken is enjoying life. He continues to be monitored and has been able to have phone consultations every three to six months throughout the COVID-19 pandemic.

More information: Gerhardt Attard et al, Abiraterone acetate and prednisolone with or without enzalutamide for high-risk non-metastatic prostate cancer: a meta-analysis of primary results from two randomised controlled phase 3 trials of the STAMPEDE platform protocol, *The Lancet* (2021). [DOI: 10.1016/S0140-6736\(21\)02437-5](https://doi.org/10.1016/S0140-6736(21)02437-5)

Provided by Institute of Cancer Research

Citation: Abiraterone could halve risk of prostate cancer death for some patients (2021, December 28) retrieved 23 November 2023 from <https://medicalxpress.com/news/2021-12-abiraterone-halve-prostate-cancer-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.