


Three-dose hepatitis B vaccine regimen protects people with HIV

October 20 2022



Hepatitis B virus particles (red). Credit: CDC/NIAID

A three-dose course of the hepatitis B vaccine HEPLISAV-B has fully protected adults living with HIV who had never been vaccinated against or infected with the hepatitis B virus (HBV), according to study findings



presented today at the IDWeek conference in Washington, D.C.

HBV is spread primarily through sexual contact and sharing of contaminated needles. It causes chronic hepatitis B infection and can lead to progressive liver disease. People living with HIV, including those who are taking antiretroviral therapy, are at greater risk of liver-related illness and death when co-infected with HBV.

According to the Centers for Disease Control and Prevention, 10% of U.S. adults with HIV also have hepatitis B. People living with HIV are less likely to produce a protective immunological response to HBV vaccination. The HEPLISAV-B <u>vaccine</u>, made by the Dynavax Technologies Corp. based in Emeryville, California, was approved in 2017 by the U.S. Food and Drug Administration as a two-dose vaccine regimen for adults. However, little was known about its protective effects in people living with HIV.

Led by study chairs Kenneth E. Sherman, M.D., Ph.D., of the University of Cincinnati College of Medicine, and Kristen Marks, M.D., of Weill Cornell Medical College, New York City, the researchers tested a three-dose course of HEPLISAV-B among 68 adults living with HIV at 38 sites in the United States, South Africa, and Thailand. None of the participants had received a previous HBV vaccination or had evidence of a previous HBV infection; all were on antiretroviral therapy.

Following the initial dose of HEPLISAV-B vaccine 0.5 milliliter (mL) as an <u>intramuscular injection</u>, study participants received additional doses at four weeks and 24 weeks. The objective of this part of the study was to assess anti-HBV surface antibodies (HBsAbs) greater than or equal to 10 milli-international units per milliliter (mIU/mL) at week 28 (defined as seroprotection from HBV) and to assess the vaccine's safety.

All participants achieved seroprotection with 88% of participants



achieving HbsAb levels greater than 1000 mIU/mL, according to findings presented today. High <u>antibody levels</u> are thought to be associated with long-term vaccine durability. At eight weeks after the second dose, 94.4% of participants achieved seroprotection; this percentage increased to 98.5% by week 24 prior to the third dose. The most common side effects related to vaccination were injection site pain, malaise, fatigue, muscle aches and headaches.

The international study will continue to examine the effects of two-dose HEPLISAV-B, as well as a three-dose regimen of another hepatitis B vaccine (ENGERIX-B, manufactured by GSK) among adult participants with HIV who were previously vaccinated against HBV but who did not achieve an adequate immunologic response. The clinical trial is estimated to complete vaccinations in March 2023.

Provided by NIH/National Institute of Allergy and Infectious Diseases

Citation: Three-dose hepatitis B vaccine regimen protects people with HIV (2022, October 20) retrieved 5 July 2023 from https://medicalxpress.com/news/2022-10-three-dose-hepatitis-vaccine-regimen-people.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.