

Yes, adults can get RSV and it can be severe

November 4 2022, by Dipali Pathak



Credit: Unsplash/CC0 Public Domain

Respiratory syncytial virus, or RSV, is a respiratory virus that is a common cause of lower respiratory illness and is most recognized for causing annual epidemics that lead to an increase in hospitalizations of children in their first or second year of life. However, it's a common misconception that RSV is a mild respiratory illness in adults when, in

fact, it can lead to symptoms as serious as those of influenza. A Baylor College of Medicine infectious diseases expert explains.

According to Dr. Robert Atmar, professor of infectious diseases at Baylor, the consequences of lower respiratory tract illnesses such as RSV can be severe, particularly for adults over 65 and those who have chronic underlying lung disease, [heart disease](#) or diabetes.

"One of the reasons that persons with these diseases are at higher risk is because of the underlying disease—the function of the heart, lungs and [immune system](#) are already compromised and less able to handle the stress associated with the infection," he said.

Historically RSV has been fairly predictable and usually occurs from October through March or April, but the COVID-19 pandemic disrupted this pattern. This year, physicians saw an increase of RSV cases starting at the end of September and are hoping that it's peaking now.

RSV is more commonly recognized in children because of its characteristic clinical findings such as bronchiolitis and the availability of diagnostic tests for pediatrics that are sensitive. Adults are not often tested for RSV and, according to Atmar, only in recent years have molecular diagnostics become more available for use in adults.

RSV can cause respiratory illness characterized by cough, shortness of breath, runny nose and nasal congestion. Fever is less common but can occur. It can start as an upper respiratory illness, like a cold, that moves to the chest as a chest cold associated with cough and possibly pneumonia. Wheezing also may be a symptom in adults, particularly those with a [chronic lung disease](#) such as COPD or asthma.

Contact with an infected individual is the most common way that RSV is transmitted, so handwashing is very important. Atmar also said that the

measures we took to decrease the transmission of COVID-19 also impacted the spread of other viruses, so those measures, such as wearing masks or respiratory hygiene, also could help prevent the transmission of RSV.

While there is no specific treatment for RSV, Atmar recommends symptomatic therapy (treatment of symptoms) such as taking a decongestant for nasal congestion or antitussives for a cough.

News of a potential RSV vaccine for [pregnant women](#) is currently making headlines and Atmar said that by this time next year, there also may be a vaccine option for older adults.

"This is not necessarily just a mild infection in all adults. It can be a very serious infection, particularly in [older adults](#) and those who have chronic heart and lung disease or diabetes and it's for those reasons that vaccines are being developed for these at-risk persons. If those vaccines become available it's going to be important for those groups approved to take it, much like they take the flu vaccine," Atmar said. "It's worth preventing if and when we have an effective [vaccine](#) available to do that, and we may have that within the next year or two."

Provided by Baylor College of Medicine

Citation: Yes, adults can get RSV and it can be severe (2022, November 4) retrieved 30 December 2022 from <https://medicalxpress.com/news/2022-11-adults-rsv-severe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--