


Transcatheter mitral valve repair in heart failure patients significantly reduces hospitalizations and improves survival

March 5 2023



Credit: Unsplash/CC0 Public Domain

Transcatheter mitral valve repair for heart failure patients with mitral regurgitation can reduce the long-term rate of hospitalizations by almost

50 percent, and death by nearly 30 percent, compared with heart failure patients who don't undergo the minimally invasive procedure.

These are the breakthrough findings from a new study led by a researcher from the Icahn School of Medicine at Mount Sinai. This multi-center trial is the largest trial to examine the safety and effectiveness of transcatheter [mitral-valve repair](#) in a [heart failure](#) population using Abbott's MitraClip system. It shows this [treatment option](#) significantly improves outcomes for patients with heart failure that do not respond to [conventional treatments](#).

The five-year results from the "Cardiovascular Outcomes Assessment of the MitraClip Percutaneous Device" study, or COAPT, were announced Sunday, March 5, in a Late Breaking Clinical Trial presentation at the American College of Cardiology Scientific Sessions Together with World Congress of Cardiology (ACC.23/WCC) in New Orleans, and published in *The New England Journal of Medicine*.

"Treating severe secondary [mitral regurgitation](#) in appropriate patients with cardiomyopathy is important—our study shows that five years after the MitraClip procedure, patients feel better, are hospitalized less frequently, and live longer," says lead author Gregg W. Stone, MD, Director of Academic Affairs for the Mount Sinai Health System and Professor of Medicine (Cardiology), and Population Health Science and Policy, at Icahn Mount Sinai. "It's critical for physicians to recognize mitral regurgitation in patients with cardiomyopathy, then treat this secondary issue as early as possible to improve outcomes in this heart failure group."

Roughly 30 percent of patients with left ventricular cardiomyopathy—the most common type of heart failure, in which the heart's main chamber (the left ventricle) becomes enlarged and can't properly pump blood out of the heart—develop a secondary heart

condition called severe mitral valve regurgitation. Secondary mitral valve regurgitation develops when the mitral valve, which controls the flow of blood from the [left atrium](#) into the left ventricle, becomes distorted from the enlarged [left ventricle](#) so its leaflets do not completely close. This causes blood to leak backwards, increases the pressure in the heart, and puts patients at increased risk of hospitalization and death. In most patients, this condition can be treated with a minimally invasive procedure called transcatheter edge-to-edge repair (TEER) in which the leaflets of the mitral valve are clipped together. The procedure is commonly performed with a device called the MitraClip, manufactured by Abbott.

In the COAPT study, researchers investigated whether treating the severe secondary mitral valve regurgitation with TEER, which has no direct effect on the underlying weakened heart muscle, would improve overall outcomes in [heart failure patients](#) beyond [medical therapy](#) alone. The two-year results, published in 2018, showed for the first time that treating secondary mitral valve regurgitation improved patients' symptoms, reduced hospitalizations, and led them to live longer. Their latest five-year results show further significant findings.

Researchers analyzed 614 patients enrolled between December 27, 2012, and June 23, 2017, at 78 sites in the United States and Canada. All patients had cardiomyopathy and secondary, severe mitral valve regurgitation despite treatment with optimal medical therapy for heart failure. Half of those patients continued their heart failure medication, while the other half underwent transcatheter valve repair with the MitraClip while continuing on the heart failure medication. Over the five years following treatment, the yearly rates of heart failure hospitalizations were 33.1 percent in the MitraClip group compared to 57.2 percent with patients treated with medications only, which was a 47 percent reduction. Deaths from heart failure were reduced by 29 percent in the MitraClip group compared with patients treated with medications

only, and all-cause death was reduced by 28 percent.

Even though patients in the MitraClip group had fewer deaths and hospitalizations after successful treatment, at the end of five years, 73.6 percent of them died or had one or more heart failure hospitalizations (compared to 91.5 percent of patients in the medication-only group). Dr. Stone says this result highlights the need for advanced therapies to treat these high-risk patients.

"Within five years, heart failure patients treated with MitraClip were on average alive and out of the hospital for 229 days more—nearly 8 months—than patients treated with medications alone. Thus, MitraClip treatment provided great benefits for these patients in terms of longevity and quality of life," noted Dr. Stone. "However, the TEER procedure doesn't cure the underlying damaged heart muscle, and the long-term survival of these patients is worse than for many cancers. New medical therapies and long-term device-based treatments for [heart failure](#) that can lower the pressures and blood volume within the heart or take over its pumping function are essential to further improve the quality of life and longevity in these desperate patients."

More information: Gregg W. Stone et al, Five-Year Follow-up after Transcatheter Repair of Secondary Mitral Regurgitation, *New England Journal of Medicine* (2023). [DOI: 10.1056/NEJMoa2300213](https://doi.org/10.1056/NEJMoa2300213)

Provided by The Mount Sinai Hospital

Citation: Transcatheter mitral valve repair in heart failure patients significantly reduces hospitalizations and improves survival (2023, March 5) retrieved 7 March 2023 from <https://medicalxpress.com/news/2023-03-transcatheter-mitral-valve-heart-failure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.