


Study explains how antidepressant increases brain plasticity

April 5 2023



Fluoxetine treatment promotes contextual and cued fear erasure which depends on TrkB expression in PV interneurons. **a** Mating strategy to obtain wild-type and PV-specific heterozygous TrkB knockout (PV-TrkB hCKO) mice. **b** Scheme of the fear-conditioning paradigm. Mice were conditioned by pairing a tone and an electric shock in context A (**c**), and then one group was treated with fluoxetine (24 mg/kg/day). After 2 weeks, mice were subjected to 2 days of fear extinction training: day 1 (Ext1), day 2 (Ext2) wt (**d**), PV-TrkB hCKO (**e**). After



1 week, mice were tested for spontaneous recovery (SR) in context B (\mathbf{g} , \mathbf{h}) and fear renewal (FR) in context A (\mathbf{i} , \mathbf{j}). \mathbf{c} Freezing was similarly increased during the conditioning/acquisition phase in both WT and hCKO mice and both genotypes reached a similar level of acquisition (two-way ANOVA, Conditioning, F(4, 270) = 21.94, P

Citation: Study explains how antidepressant increases brain plasticity (2023, April 5) retrieved 3 July 2023 from https://medicalxpress.com/news/2023-04-antidepressant-brain-plasticity.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.