

Stroke: Know the signs and risk factors

May 5 2023, by Laurel Kelly



Credit: CC0 Public Domain

May is National Stroke Awareness Month, which makes this a good time to learn more about the signs and risk factors for stroke.

Stroke is one of the leading causes of death in the U.S. and a major

cause of serious disability for adults. More than 795,000 people in the U.S. have a stroke each year, according to the Centers for Disease Control and Prevention.

A stroke occurs when the [blood supply](#) to part of your brain is interrupted or reduced, preventing [brain tissue](#) from getting oxygen and nutrients. Brain cells begin to die in minutes.

A stroke is a [medical emergency](#), and prompt treatment is crucial. Early action can reduce [brain damage](#) and other complications. If you or someone you know is experiencing a stroke, you should call 911 and seek emergency medical care right away.

Signs of stroke

To recognize the signs of stroke, remember the acronym FAST:

- **Face:** Does the face droop on one side when the person tries to smile?
- **Arms:** Is one arm lower when the person tries to raise both arms?
- **Speech:** Can the person repeat a simple sentence? Is speech slurred or hard to understand?
- **Time:** During a stroke, every minute counts. If you observe any of these signs, call 911 or your local emergency number immediately.

Risk factors

Many factors can increase the risk of stroke, including:

- **Age:** People 55 or older have a higher risk of stroke than [younger people](#).
- **Race or ethnicity:** People who are African American and

Hispanic have a higher risk of stroke than people of other races or ethnicities.

- Sex: Men have a higher risk of stroke than women. Women are usually older when they have strokes, and they're more likely to die of strokes than men, however.
- Hormones: Use of birth control pills or hormone therapies that include estrogen increases risk.

Potentially treatable stroke risk factors include lifestyle and medical factors.

Lifestyle risk factors include:

- Being overweight or obese
- Physical inactivity
- Heavy or binge drinking
- Use of illegal drugs, such as cocaine and methamphetamine

Medical risk factors include:

- High blood pressure
- Cigarette smoking or secondhand smoke exposure
- High cholesterol
- Diabetes
- Obstructive sleep apnea
- Cardiovascular disease, including [heart failure](#), heart defects, heart infection or irregular heart rhythm
- Personal or family history of stroke, heart attack or transient ischemic attack
- COVID-19 infection

Stroke rehabilitation

A stroke can cause temporary or permanent disabilities, depending on how long the brain lacks blood flow and which part was affected. Complications can include paralysis or loss of muscle movement, difficulty talking or swallowing, [memory loss](#) or thinking difficulties, emotional problems, pain, and changes in behavior and self-care ability.

Stroke rehabilitation is an important part of recovery after stroke. There are many approaches to stroke rehabilitation. Your rehabilitation plan will depend on the part of the body or type of ability affected by your [stroke](#).

©2023 Mayo Clinic News Network.

Distributed by Tribune Content Agency, LLC.

Citation: Stroke: Know the signs and risk factors (2023, May 5) retrieved 8 May 2023 from <https://medicalxpress.com/news/2023-05-factors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--