


Targeting telomeres could be an effective therapeutic strategy against lung cancer, according to study

May 25 2023



Increased amplification frequency, copy number values and mRNA expression of TERT in NSCLC patients, and reduced tumor implantation in TERT-deficient mice upon lung tumor induction. Amplification frequency (A), copy number values (B) mRNA expression levels of *TERT* (C) and Pearson correlation of mRNA expression with copy number values of *TERT* (D) in lung tissues from NSCLC patients, and survival probability in NSCLC patients with high and low

TERT expression (**E**) obtained from the Kaplan–Meier Plotter database. **F** Correlation between the expression of *TERT* and immune infiltrates in NSCLC patients from the TCGA using the TIMER 2.0 database. **G, H** Generation of *Tert*^{+/+} and G3 *Tert*^{-/-} mice and protocol for the induction of Lewis Lung Carcinoma (LLC). **G** Heterozygous *Tert*^{+/-} mice were crossed to obtain *Tert*^{+/+} and G1 *Tert*^{-/-} mice, and successive crosses between G1 *Tert*^{-/-} and then G2 *Tert*^{-/-} were set to generate G3 *Tert*^{-/-} mice. **H** 1 × 10⁶ Lewis cells suspended in 100 µl of DMEM or equal volume of DMEM (controls) were injected via the tail vein of 10–12 weeks old *Tert*^{+/+} and G3 *Tert*^{-/-} male mice on day 0 (D0). An in vivo follow follow-up of survival was carried out until sample collection on day 14 (D14). Kaplan–Meier survival curves (**I**), representative images of LLC-challenged *Tert*^{+/+} and G3 *Tert*^{-/-} lungs and controls (H&E) (**J**), and quantification of lung tumor area (**K, L**) and foci (**M**) in *Tert*^{+/+} and G3 *Tert*^{-/-} mice. **N** Representative Telomeric repeat amplification protocol (TRAP) using S-100 lung extracts from LLC-challenged *Tert*^{+/+} and G3 *Tert*^{-/-} mice and controls, where protein concentration is indicated. Extracts were treated (+) or not (-) with RNase as a negative control (exposition time: 48 h). An internal control (IC) for PCR efficiency was also included and arrows indicate the lane used for quantification. **O** Quantification of Telomerase activity in lung extracts from LLC-challenged *Tert*^{+/+} and G3 *Tert*^{-/-} mice and controls expressed in arbitrary units (a.u). **P** Lung tissue mRNA expression levels of *Tert* normalized to 18S expression in *Tert*^{+/+} and G3 *Tert*^{-/-} mice. Data are expressed as mean ± SEM (the number of mice is indicated in each case). ****p*

Citation: Targeting telomeres could be an effective therapeutic strategy against lung cancer, according to study (2023, May 25) retrieved 4 July 2023 from <https://medicalxpress.com/news/2023-05-telomeres-effective-therapeutic-strategy-lung.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.