

Gaining ground on sickle cell disease

15 July 2008

Although sickle cell disease is a single-gene disorder, its symptoms are highly variable. In a study published online July 14 by the *Proceedings of the National Academy of Sciences*, scientists at Children's Hospital Boston and the Dana Farber Cancer Institute (DFCI), in collaboration with the Broad Institute of MIT and Harvard, report five gene variants that could potentially be helpful in predicting sickle cell disease severity, perhaps even leading to better treatment approaches in the future.

The gene variants influence blood levels of fetal hemoglobin (HbF), which are known to affect symptom severity in sickle cell disease—with some patients experiencing frequent, severe pain crises and organ damage, while others are scarcely aware of their disease.

"Our study is a first step towards a better understanding of fetal hemoglobin regulation in patients with sickle cell disease," says Guillaume Lettre, PhD, of the Broad Institute and Children's Hospital Boston, and co-first author on the paper. "But further validation experiments are needed before these findings can become useful in the clinic."

"Eventually, understanding the factors giving rise to heterogeneity in HbF levels might allow us to take severely affected patients and make them more like those with more benign symptoms," adds Vijay Sankaran, co-first author on the paper with Lettre and an MD-PhD student in the laboratory of Stuart Orkin, MD. (Orkin is chair of pediatric oncology at DFCI and a Howard Hughes Medical Institute investigator at Children's.)

In sickle cell disease, a single genetic mutation results in the production of an abnormal type of hemoglobin, the main component of red blood cells. The abnormal hemoglobin molecules tend to form long chains, causing red blood cells to become stiff and sickle-shaped. The distorted cells have difficulty passing through blood vessels and can block the smaller vessels, resulting in severe

pain and eventual organ damage as tissues are robbed of their blood supply. The sickle-shaped red blood cells also have a very short lifespan, causing patients to be chronically anemic.

Previous research had established that retaining high levels of another type of hemoglobin—HbF, found at high levels in the fetus—can ameliorate sickle cell disease symptoms. At birth, HbF comprises between 50 to 95 percent of a child's hemoglobin, gradually declining as the switch is made to adult hemoglobin production -- consistent with clinicians' observations that newborns diagnosed with sickle cell disease usually do not become symptomatic until they are about a year old. Population studies in Saudi Arabia and parts of India had identified groups of sickle cell patients with very high levels of HbF and relatively benign forms of the disease, and additional epidemiologic studies led by Orah Platt, MD, chief of laboratory medicine at Children's, showed that HbF is an ameliorating factor. "The more you have, the better off you are," says Sankaran.

Studying 1600 patients with sickle cell disease, the researchers found that previously identified DNA sequence variants in three chromosome locations (small regions on chromosome 2, 6, and 11) were associated with high or low HbF levels. When they added these five variants to a model previously designed by Platt to predict disease severity, which also factors in age, sex, degree of anemia and HbF levels, the model's predictive ability was enhanced.

The findings need to be validated in large, prospective clinical studies, but the researchers are hopeful about the possible future clinical implications of their work. "As we find gene variants that regulate HbF levels or predict severity, we might eventually want to genotype patients for these variants, to get more predictive information on their disease," Sankaran says.

Finally, once this study is validated, understanding how these variants actually affect HbF levels might someday lead to new drugs that do the same thing.

"If we can gain better insight into what these variants are doing, we may eventually have better, more targeted therapies for sickle cell disease," adds Sankaran.

Source: Children's Hospital Boston

APA citation: Gaining ground on sickle cell disease (2008, July 15) retrieved 11 June 2021 from <https://medicalxpress.com/news/2008-07-gaining-ground-sickle-cell-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.