

Survey identifies factors associated with early detection of melanoma in older men

20 April 2009

Older men whose melanoma is detected by a physician are more likely to have thinner and therefore more treatable tumors at diagnosis, according to results of a survey published in the April issue of *Archives of Dermatology*, one of the JAMA/Archives journals. A second analysis of the same survey data finds that physician detection of thin melanoma is more common in those who are 65 or older, have cancers on their backs or who have a history of atypical moles.

Melanoma is becoming more common and mortality rates from melanoma are steadily increasing among older men, according to background information in one of the articles. Tumor thickness at diagnosis strongly predicts the management and outcomes of melanoma, and the thickest tumors (4 millimeters or thicker) are increasingly common in white men age 60 and older. "Rigorous assessment of behavioral, social and medical access factors that differ between men 40 years or older with thinner vs. thicker melanomas may identify potential modifiable variables," the authors write. "A clearer understanding of these factors provides fundamental knowledge for additional studies and public health messages aimed at earlier melanoma detection in this high-risk subset of men."

In one study, Susan M. Swetter, M.D., of Stanford University Medical Center, Calif., and colleagues surveyed 227 men age 40 and older between 2004 and 2006, within three months of their melanoma diagnosis. The men responded to questions about their previous melanoma awareness, skin examination practices, how their cancer was discovered and social and medical care factors.

Of the 227 men, 57 (25.1 percent) had tumors thicker than 2 millimeters. These men were more likely to have less than a high school education, less likely to have a history of atypical nevi (moles) and their melanomas were more often discovered by the patient themselves or a friend or family

member than by a physician. Men with thinner melanomas were more likely to have previous knowledge of melanoma, have paid attention to skin cancer detection information, be interested in health topics and be aware of the importance of physician skin examination. Tumor thickness was not associated with patients' age, whether or not they were married or lived with a partner, skin cancer history, sun sensitivity or the anatomic location of the cancer.

Overall, few patients were aware of melanoma warning signs (less than 20 percent), practiced skin self-examination (less than 50 percent) or used the Internet (less than 14 percent) as a source of skin cancer information.

"For men 40 years or older, who constitute more than half of all melanoma deaths in the United States, we have identified at least two key variables (physician skin examination and improved public awareness, particularly for patients in lower socioeconomic groups) as major targets for new interventions to promote earlier melanoma detection," the authors write. "Public education, in particular targeting less-educated, middle-aged and older men for improved self-examination and physician skin surveillance, should become an integral component of skin cancer risk reduction strategies promoted by cancer advocacy organizations."

In another study, Alan C. Geller, M.P.H., R.N., of Boston University, and colleagues report a separate analysis of the same survey data. Patients whose melanomas were detected by a physician tended to be older—57 percent were 65 years or older, compared with 42 percent of patients who detected cancer themselves and 34 percent whose cancer was detected by another non-physician. This may be because older patients visit their physicians more frequently, providing more opportunities for skin examination, and they may rely more heavily on physician skin evaluation

because they tend to have poorer eyesight and are less likely to have a partner.

Forty-six percent of physician-detected melanomas, 16 percent of self-detected melanomas and 57 percent of melanomas detected by other means were on the patient's back; these back-of-the-body melanomas were smaller than 2 millimeters in 92 percent of physician-detected cases, 63 percent of self-detected cases and 76 percent of those detected by other means.

"Skin screenings of at-risk middle-aged and older American men can be integrated into the routine physical examination, with particular emphasis on hard-to-see areas, such as the back of the body," the authors conclude. "'Watch your back' professional education campaigns should be promoted by [skin cancer](#) advocacy organizations and should incorporate the importance of physician screening and the benefit of spouse or partner assistance for early detection of [melanoma](#), particularly in the high-risk population of middle-aged and older men."

More information: Arch Dermatol. 2009;145[4]:397-404, 409-414.

Source: JAMA and Archives Journals ([news](#) : [web](#))

APA citation: Survey identifies factors associated with early detection of melanoma in older men (2009, April 20) retrieved 20 June 2022 from <https://medicalxpress.com/news/2009-04-survey-factors-early-melanoma-older.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.