

Infiltrating blood-derived macrophages play a role in recovery from spinal cord injury in mice

July 28 2009

Although macrophages are known as essential players in wound healing, their contribution to recovery from spinal cord injury is a subject of debate.

Using a mouse model of spinal injury, Michal Schwartz and colleagues from the Weizmann institute of Science, Rehovot, Israel tested the effect of macrophages on the recovery process after injury and demonstrate an important anti-inflammatory role for a subset of infiltrating monocyte-derived macrophages that is dependent upon their expression of the anti-inflammatory molecule interleukin-10.

These results suggest that this subset of macrophages may have a beneficial effect on [spinal cord](#) injuries.

More information: Shechter R, London A, Varol C, Raposo C, Cusimano M, et al. (2009) Infiltrating Blood-Derived Macrophages Are Vital [Cells](#) Playing an Anti-inflammatory Role in Recovery from Spinal Cord Injury in Mice. *PLoS Med* 6(7): e1000113.

[doi:10.1371/journal.pmed.1000113](https://doi.org/10.1371/journal.pmed.1000113), medicine.plosjournals.org/perldoc/document&doi=1000113

Source: Public Library of Science ([news](#) : [web](#))

Citation: Infiltrating blood-derived macrophages play a role in recovery from spinal cord injury in mice (2009, July 28) retrieved 1 February 2024 from <https://medicalxpress.com/news/2009-07-infiltrating-blood-derived-macrophages-role-recovery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.