

# Short-term school closures may worsen flu pandemics, study finds

30 December 2009

Closing schools for less than two weeks during a flu pandemic may increase infection rates and prolong an epidemic, say University of Pittsburgh researchers in a study published ahead-of-print and online in the *Journal of Public Health Management and Practice*. The findings, developed from a series of computer simulations based on U.S. census data, indicate that schools may need to be closed for at least eight weeks in order to significantly decrease the spread of infection.

The value of school closures has been debated as a possible strategy to stem or slow the current H1N1 [influenza pandemic](#). Indeed, hundreds of schools across the country have been closed at different periods during 2009 for fear the virus would spread more quickly if they stayed open.

"Although closing schools may seem like a reasonable way to slow the spread of flu, we found that it was not effective unless sustained for at least eight weeks after implementation," said study lead author, Bruce Lee, M.D., M.B.A., assistant professor of medicine, epidemiology and biomedical informatics, University of Pittsburgh. Closing schools quickly at the start of an outbreak was much less important than keeping them closed continually throughout the epidemic, he added.

According to study authors, short-duration school closures can increase transmission rates by returning susceptible students back to school in the middle of an epidemic when they are most vulnerable to infection.

The study also found that identifying sick students individually and keeping them from attending school had minimal impact on an epidemic. In addition, there were no significant differences between individual school closures and system-wide closures in mitigating an epidemic.

The study was based on an agent-based computer

simulation model of Allegheny County, Pa., that represented the county's population, school systems, workplaces, households and communities. Simulations were based on the movement of residents each weekday from their households to designated workplaces or schools, and included 1.2 million people?200,000 of whom were school-aged children. The study also included more than 500,000 households and nearly 300 schools.

Provided by University of Pittsburgh Schools of the Health Sciences

APA citation: Short-term school closures may worsen flu pandemics, study finds (2009, December 30) retrieved 24 September 2022 from <https://medicalxpress.com/news/2009-12-short-term-school-closures-worsen-flu.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*