

New SBI, ACR recommendations suggest breast cancer screening should begin at age 40

4 January 2010

The new recommendations from the Society of Breast Imaging (SBI) and the American College of Radiology (ACR) on breast cancer screening, published in the January issue of the *Journal of the American College of Radiology* (JACR), state that breast cancer screening should begin at age 40 and earlier in high-risk patients. The recommendations also suggest appropriate utilization of medical imaging modalities such as mammography, magnetic resonance imaging (MRI), and ultrasound for breast cancer screening.

"The significant decrease in <u>breast cancer</u> mortality, which amounts to nearly 30 percent since 1990, is a major medical success and is due largely to earlier detection of breast cancer through <u>mammography</u> screening," said Carol H. Lee, MD. "For women with the highest risk of developing breast cancer, screening technologies in addition to mammography have been adopted," said Lee.

The new recommendations released by the SBI and ACR include recommendations for imaging screening for breast cancer by imaging technique (mammography, MRI, and ultrasound) and by risk factor. The recommendations state that the average patient should begin annual breast cancer screening at age 40. High-risk patients should begin by age 30 but not before 25. "Evidence to support the recommendation for regular periodic screening mammography comes from the results of several randomized trials (RCTs) conducted in Europe and North America that included a total of nearly 500,000 women. Overall, based on a metaanalysis of the RCTs, there was a 26 percent reduction in mortality," said Lee.

"It should be remembered that mammography is the only imaging modality that has been proven to decrease mortality from breast cancer. However major efforts continue to build on this success by

developing additional methods to screen for early breast cancer," she said.

"The SBI and ACR wish to remind women and their physicians that in those instances in which there is a concern that risk of developing cancer is considerably elevated from that of the general population, consultation with appropriate experts in breast cancer genetics and/or high risk management is desirable," said Lee.

Provided by American College of Radiology



APA citation: New SBI, ACR recommendations suggest breast cancer screening should begin at age 40 (2010, January 4) retrieved 2 May 2021 from <u>https://medicalxpress.com/news/2010-01-sbi-acr-breast-cancer-screening.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.