

Regional protocols improve quality of care for aortic dissection patients

August 10 2010

Using a standardized protocol within a regional health network, a multidisciplinary acute aortic dissection (AAD) program resulted in a 43 percent reduction in time to diagnosis for all AAD patients over the course of five years, based on study results published in the July edition of *Circulation: Cardiovascular Quality and Outcomes*, an American Heart Association journal.

While the incidence rate of AAD is rare—approximately three cases per 100,000 persons annually—these patients have an in-hospital mortality rate of 26 percent, and for those patients with type A aortic dissection, the [mortality risk](#) is 1 percent to 2 percent per hour until emergency surgical repair is performed. "It is therefore critical that AAD be recognized promptly and that surgical care be provided expeditiously," according to background information of the study.

"Aortic dissection has a much higher mortality rate than more prolific disease states, such as severe heart attacks," explained Kevin M. Harris, MD, who is co-director of the Acute Aortic Dissection Program at the Minneapolis Heart Institute® at Abbott Northwestern Hospital in Minneapolis, Minn.

The International Registry of Acute Aortic Dissection (IRAD) estimates that the average time between an emergency room arrival and the diagnosis of acute aortic dissections is 4.3 hours.

In an effort to accelerate the time between hospital arrival to diagnosis

and diagnosis to intervention, Harris and his colleagues at the Minneapolis Heart Institute® and Minneapolis Heart Institute Foundation established quality improvement protocols and standardized care measures throughout 32 community hospitals in Minnesota, North Dakota and western Wisconsin. As part of an ongoing AAD educational program, the participating facilities received more than 60 lectures and training sessions on AAD and received program-related protocols and toolkits.

During the study, the researchers examined 101 patients with aortic dissection—68 percent of whom had aortic dissection affecting the ascending aorta (Type A) and 76 percent were transferred to the aortic dissection center from community hospitals.

Overall, the study showed that the system-wide education and coordination efforts resulted in a 43 percent reduction in time to diagnosis for all cases of AAD and a 55 percent reduction in the time from diagnosis to intervention in surgically managed type A patients.

"The time was reduced from 279 minutes to 160 minutes, which equates to a two-hour time savings," said Dr. Harris.

The researchers also reported a 30 percent decrease between the time a patient was diagnosed with aortic dissection and the time they presented to the operating room. "The times prior to use of the protocols were 728 minutes compared to 360 minutes after implementation of the protocols, and we've cut the time of transfer from arrival to the operating room basically in half," said Dr. Harris.

The study authors noted that no new technologies or significant financial costs were required to implement the program. "This systematic approach to AAD care represents a new paradigm for the treatment of this life-threatening condition," they concluded.

Provided by Minneapolis Heart Institute Foundation

Citation: Regional protocols improve quality of care for aortic dissection patients (2010, August 10) retrieved 7 March 2023 from <https://medicalxpress.com/news/2010-08-regional-protocols-quality-aortic-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.