

Room light before bedtime may impact sleep quality, blood pressure and diabetes risk

13 January 2011

According to a recent study accepted for publication in The Endocrine Society's *Journal of Clinical Endocrinology & Metabolism (JCEM)*, exposure to electrical light between dusk and bedtime strongly suppresses melatonin levels and may impact physiologic processes regulated by melatonin signaling, such as sleepiness, thermoregulation, blood pressure and glucose homeostasis.

Melatonin is a hormone produced at night by the pineal gland in the brain. In addition to its role in regulating the sleep-wake cycle, melatonin has been shown to lower [blood pressure](#) and body temperature and has also been explored as a treatment option for insomnia, hypertension and cancer. In modern society, people are routinely exposed to electrical lighting during evening hours to partake in work, recreational and social activities. This study sought to understand whether exposure to room light in the late evening may inhibit melatonin production.

"On a daily basis, millions of people choose to keep the lights on prior to bedtime and during the usual hours of sleep," said Joshua Gooley, PhD, of Brigham and Women's Hospital and Harvard Medical School in Boston, Mass. and lead author of the study. "Our study shows that this exposure to indoor light has a strong suppressive effect on the hormone melatonin. This could, in turn, have effects on [sleep quality](#) and the body's ability to regulate body temperature, blood pressure and glucose levels."

In this study, researchers evaluated 116 healthy volunteers aged 18-30 years who were exposed to room light or dim light in the eight hours preceding bedtime for five consecutive days. An intravenous catheter was inserted into the forearms of study participants for continuous collection of blood

plasma every 30-60 minutes for melatonin measurements. Results showed exposure to room light before [bedtime](#) shortened melatonin duration by about 90 minutes when compared to dim light exposure. Furthermore, exposure to room light during the usual hours of sleep suppressed melatonin by greater than 50 percent.

"Given that chronic light suppression of melatonin has been hypothesized to increase relative risk for some types of cancer and that melatonin receptor genes have been linked to type 2 diabetes, our findings could have important health implications for shift workers who are exposed to indoor light at night over the course of many years," said Gooley. "Further research is still needed to both substantiate melatonin suppression as a significant risk factor for breast cancer and determine the mechanisms by which [melatonin](#) regulates glucose metabolism."

More information: The article, "Exposure to room light prior to bedtime suppresses melatonin onset and shortens melatonin duration in humans," appears in the March 2011 issue of *JCEM*.

Provided by The Endocrine Society

APA citation: Room light before bedtime may impact sleep quality, blood pressure and diabetes risk (2011, January 13) retrieved 25 May 2022 from <https://medicalxpress.com/news/2011-01-room-bedtime-impact-quality-blood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.