

Study finds preterm birth clinic attendance leads to major reduction in infant disability

11 February 2011

In a study to be presented today at the Society for Maternal-Fetal Medicine's (SMFM) annual meeting, The Pregnancy Meeting, in San Francisco, researchers will present findings that show that when women at high risk for preterm birth participated in a preterm birth prevention clinic, more women delivered full term babies and there were fewer cases of infant morbidity.

The National Center for Health Statistic reports that in 2008, 12.3% of babies were born prematurely. Women who have had a prior preterm birth are at high risk to have another one. In 2008, Intermountain Healthcare created a preterm birth (PTB) prevention clinic to focus care for this high-risk population.

"We wanted to take a very aggressive approach to treating women with a history of preterm birth," said Sean Esplin, M.D., of Intermountain Healthcare and one of the studies authors. "We gathered together the best treatments for women at high risk for preterm labor and administered them in a systematic way." He continued, "Then we designed a study to see if the intervention leads to better results in future pregnancies."

Esplin and his colleagues conducted a retrospective review of women with a single, non-anomalous fetus and 21 documented previous spontaneous PTB

Two hundred and thirty-two patients (70 PTB Prevention Clinic and 162 Usual-Care patients) met inclusion criteria. Groups had similar previous [pregnancy](#) characteristics. PTB Prevention Clinic patients had increased utilization of resources (including more cervical length ultrasounds and higher rates of use of prophylactic 17OHPC) and delivered at later gestational ages. Rates of NICU admission were similar between groups (44.3% vs. 40.7%, $p=0.62$). However, rates of major neonatal morbidity (diagnosis of NEC, BPD, IVH, sepsis, or death) were lower among PTB Prevention Clinic

neonates (5.3% vs. 15.4%, $p=0.025$).

The study showed that among this high-risk population, referral to a consultative PTB Prevention Clinic (with standardized counseling, management recommendations, and close surveillance) resulted in a reduction in the rate of recurrent PTB prior to 37 weeks, lead to an average of a one week longer pregnancy, and reduced the rates of major neonatal morbidity.

"The study showed that participants in the preterm birth prevention clinic had a 28 percent reduction in the risk of recurrent spontaneous preterm birth, as well as reductions in infant complications and short term disabilities," said Tracy Manuck, M.D., one of the study's authors. "These are significant improvements and should lead more medical facilities to think about creating similar programs."

Provided by Society for Maternal-Fetal Medicine

APA citation: Study finds preterm birth clinic attendance leads to major reduction in infant disability (2011, February 11) retrieved 25 May 2021 from <https://medicalxpress.com/news/2011-02-preterm-birth-clinic-major-reduction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.