

Study finds some patients with cerebral palsy have asymmetric pelvic bones

10 March 2011

Johns Hopkins Children's Center researchers have discovered that most children with severe cerebral palsy have starkly asymmetric pelvic bones. The newly identified misalignment can affect how surgeries of the pelvis, spine and surrounding structures are performed, the researchers say.

The study will be published online on March 10 in the *Journal of Pediatric Orthopaedics*.

Previous studies of patients with cerebral palsy have reported asymmetry above the [pelvis](#) and misalignment of the hips, but this new report, the researchers say, is the first one to show misalignment between the two sides of the pelvic bone itself.

Most children with severe cerebral palsy have significant spinal curvatures (scoliosis) that often require surgery. Because the pelvis and the spine are connected, any surgical procedures to correct scoliosis should take into account the possibility of a misaligned pelvis, the investigators say. The degree of the asymmetry, they add, should dictate the size, type and placement of the surgical screws and rods used to stabilize the spine and pelvis in such corrective procedures.

"Surgeons preparing to operate on children with cerebral palsy should look out for pelvic asymmetry and tweak their surgical technique accordingly to achieve better outcomes and more lasting benefits," says senior investigator Paul Sponseller, M.D., chief of pediatric orthopedics at Hopkins Children's.

While performing surgeries to correct scoliosis, Sponseller started noticing a recurrent feature among his patients with severe cerebral palsy - a pronounced asymmetry between the left and right plates of the pelvic bone.

To quantify the problem, Sponseller performed three-dimensional CT scans on all of his cerebral

palsy patients undergoing scoliosis surgery over one year. All 27 patients had asymmetric pelvises with misalignment of the pelvic bones was greater than 10 degrees. Comparing these images with pelvic-bone scans of children without cerebral palsy, the researchers noted that all of them had either no misalignment or only mild asymmetry of less than 10 degrees.

Twenty-three of the 27 children (85 percent) with [cerebral palsy](#) also had windswept hips, a hallmark feature in CP patients marked by one hip facing outward and the other one rotated inward. Children with windswept hips had more pronounced pelvic asymmetry than children without windswept hips, the researchers found.

Co-investigators on the research included Phebe Ko, B.S., Paul Jameson II, B.S., and Tai-Li Chang, M.D., all of Hopkins.

Provided by Johns Hopkins Medical Institutions

APA citation: Study finds some patients with cerebral palsy have asymmetric pelvic bones (2011, March 10) retrieved 23 July 2022 from <https://medicalxpress.com/news/2011-03-patients-cerebral-palsy-asymmetric-pelvic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.