

Combination ACE inhibitor therapy increases risk of kidney failure and death

21 March 2011

Elderly patients prescribed combination angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARB) had a higher risk of kidney failure and death, according to a study published in *CMAJ (Canadian Medical Association Journal)*.

did not have the trial-proven indications and that it was frequently stopped after only a few months, even when hyperkalemia or renal dysfunction did not occur," conclude the researchers.

This study, by researchers from the University of Alberta and the University of Calgary, sought to determine the safety of combination therapy of ACE inhibitors and ARB in the clinical setting as some randomized trials indicate an increased risk of [kidney failure](#). Randomized trials may over or underestimate the risk of adverse events possibly because of patient selection bias, higher drug doses and increased monitoring.

Provided by Canadian Medical Association Journal

The researchers looked at 32 312 seniors in Alberta, Canada, aged 65 and older who were prescribed an [ACE inhibitor](#) and/or an ARB. They compared patients receiving both drugs together with patients who received only one of the drugs. They found a higher risk of adverse events such as high creatinine levels, end-stage renal disease and death in people taking combination therapy.

"We found that less than one-seventh of the elderly residents of Alberta who were given combination therapy in clinical practice had either of the conditions for which this therapy has been proven beneficial in randomized trials (i.e., proteinuria or symptomatic left ventricular systolic dysfunction despite treatment with and ACE inhibitor or an angiotensin-receptor blocker alone," writes Dr. Finlay McAlister, University of Alberta, with coauthors.

As well, they observed that within three months, most patients stopped the combination therapy. The authors speculate this may have been due to [low blood pressure](#).

"Our most striking findings were that [combination therapy](#) was commonly prescribed for patients who

APA citation: Combination ACE inhibitor therapy increases risk of kidney failure and death (2011, March 21) retrieved 13 August 2022 from <https://medicalxpress.com/news/2011-03-combination-ace-inhibitor-therapy-kidney.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.