

# Vitamin D deficiency in pneumonia patients associated with increased mortality

May 10 2011

---

A new study published in the journal *Respirology* reveals that adult patients admitted to the hospital with pneumonia are more likely to die if they have Vitamin D deficiency.

Vitamin D is known to be involved in the [innate immune response](#) to infection.

The team of researchers at Waikato Hospital and the Universities of Waikato and Otago, measured vitamin D in the blood samples of 112 adult patients admitted with community acquired pneumonia during the winter at the only acute-care hospital in Hamilton, New Zealand.

The researchers found that Vitamin D deficiency was associated with higher mortality within the first 30 days after [hospital admission](#) for pneumonia. The association between [vitamin D deficiency](#) was not explained by patient age, sex, comorbidities, the severity of the [systemic inflammatory response](#), or other known prognostic factors.

The authors conclude that "improved understanding of Vitamin D and its role in immunity may lead to better ways to prevent and/or treat pneumonia. We now need to investigate whether Vitamin D supplements could be a useful addition to pneumonia treatment and whether using supplements could help to prevent or reduce the severity of pneumonia among high-risk populations."

Provided by Wiley

Citation: Vitamin D deficiency in pneumonia patients associated with increased mortality (2011, May 10) retrieved 28 January 2023 from <https://medicalxpress.com/news/2011-05-vitamin-d-deficiency-pneumonia-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.