

Patients' brains may adapt to ADHD medication

2 February 2012



(Medical Xpress) -- New research reveals how the brain appears to adapt to compensate for the effects of long-term ADHD medication, suggesting why ADHD medication is more effective short-term than it is long-term. The study, from the Institute of Psychiatry (IoP) at King's College London is published today in the *American Journal of Psychiatry*.

Dr. Paolo Fusar-Poli and Professor Katya Rubia at the IoP at King's led the research. Prof Rubia says: "There is currently no evidence for the long-term effectiveness of stimulant medication. In fact, there is evidence that the effect of medication diminishes over time and we know from clinicians that medication doses often need to be increased over time to be as effective as they were initially. Our findings could help explain why stimulants work very well in the short term but not so well long-term."

ADHD has been associated with abnormalities in the [dopamine](#) system. Dopamine is a chemical important for motivation and attention, and is present at abnormally low levels in [patients](#) with ADHD. Stimulants, such as Ritalin, block dopamine transporters (DATs) which are responsible for the uptake of dopamine into the brain. Therefore, administering stimulants results in less dopamine being taken up by the brain and more free

dopamine being available.

However, measuring the exact levels of DAT in the brains of patients with ADHD has until now, revealed contradictory results. Early Positron Emission Tomography (PET) and Single Photon Emission Tomography (SPECT) scans found that DAT levels are elevated in the basal ganglia in ADHD patients, while later studies found them to be reduced.

The researchers aimed to explain these conflicting findings by performing a combined analysis of nine SPECT and PET studies investigating levels of DAT in a total of 169 ADHD patients compared to 173 healthy controls - out of the nine studies, only one focused on children while the others looked at adults.

They found that the level of DAT in patients with ADHD was strongly linked to whether patients had received long-term medication or not. Patients who had never taken stimulant medication had abnormally reduced DAT levels in their brain, whilst patients who had received long-term medication had abnormally elevated DAT levels.

The findings suggest that, when treated with stimulants, the [brain](#) adapts to compensate for the high levels of dopamine by building up more DATs to eliminate the abnormally high levels of dopamine. The findings therefore show that increased DAT levels in [ADHD](#) is a consequence of long-term medication rather than the disorder itself.

The authors add that further long-term studies are needed to investigate the long-term effect of this neurobiological adaptation.

More information: Fusar-Poli, P. et al. 'Striatal dopamine transporter alteration in ADHD: pathology of adaptation to psychostimulants? A meta-analysis' (1st February 2012) *American Journal of Psychiatry*

doi: 10.1176/appi.ajp.2011.11060940

Provided by King's College London

APA citation: Patients' brains may adapt to ADHD medication (2012, February 2) retrieved 3 May 2021 from <https://medicalxpress.com/news/2012-02-patients-brains-adhd-medication.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.