

44 percent of postmenopausal women with distal radius fracture have low levels of vitamin D

February 7 2012

Wrist fractures, also called distal radius fractures (DRF), are among the most common osteoporosis-related fractures occurring on average 15 years earlier than hip fractures. As vitamin D deficiency has recently been linked with muscle weakness, increased fall risks, and bone fractures, investigators sought to determine the prevalence of vitamin D deficiency among post menopausal women with DRF. The study, "Hypovitaminosis D in Postmenopausal Women with a Distal Radius Fracture," was presented today at the 2012 Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS).

Medical records of 104 post menopausal women treated for a DRF, and 107 age-matched control patients with soft tissue disease, were reviewed. Mean vitamin D levels were "significantly" lower in the DRF group of patients. Specifically, 26 percent of the DRF patients were vitamin D insufficient (having vitamin D serum levels between 20 and 32 ng/ml), and 18 percent, deficient ([serum levels](#) below 20 ng/ml), compared to 11 percent and 2 percent of patients being vitamin D insufficient and deficient, respectively, in the control group.

Further research may determine whether vitamin D supplementation (or, maintaining adequate vitamin D levels) can help prevent distal radius fractures, or prevent future fractures in patients that experience their first distal radius fracture.

Provided by American Academy of Orthopaedic Surgeons

Citation: 44 percent of postmenopausal women with distal radius fracture have low levels of vitamin D (2012, February 7) retrieved 7 May 2023 from

<https://medicalxpress.com/news/2012-02-percent-postmenopausal-women-distal-radius.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.