

# Mental stress may be harder on women's hearts

26 April 2012

Coronary artery disease continues to be a major cause of death in the U.S., killing hundreds of thousands of people per year. However, this disease burden isn't evenly divided between the sexes; significantly more men than women are diagnosed with coronary artery disease each year. The reasons behind this difference aren't well defined. Though some studies have shown that men's hearts become more constricted than women's during exercise, letting less blood flow through, women are more likely than men to have symptoms of heart trouble after emotional upsets.

Searching for the reasons behind these disparities, Charity L. Sauder, Alison E. Thompson, Terrell Myers, and Chester A. Ray, all of Penn State College of Medicine, investigated the effects of [mental stress](#) on [blood flow](#) through the heart. Their findings show that coronary blood flow actually increases in men during mental stress, but shows no change in women. These results may explain why women could be more susceptible to adverse [cardiac events](#) when under stress.

An abstract of their study entitled, "Effect of Mental Stress on Coronary Blood Flow in Humans," will be discussed at the meeting [Experimental Biology 2012](#), being held April 21-25 at the San Diego Convention Center. The abstract is sponsored by the [American Physiological Society](#) (APS), one of six scientific societies sponsoring the conference which last year attracted some 14,000 attendees.

## Stressing the Sexes

The researchers recruited 17 healthy adults, a near equal mix of men and women. Each volunteer had his or her heart rate and blood pressure measured at rest, as well as coronary vascular [conductance](#), a [Doppler ultrasound](#) measure of blood flow through the coronary blood vessels of the heart.

These volunteers then underwent the same tests while participating in three minutes of mental

arithmetic, in which the researchers had them sequentially subtract 7 starting with a random number. To increase the stress load, researchers lightly badgered the volunteers during the task, urging them to hurry up or telling them they were wrong even when they gave the correct number. At the end of the task, they underwent the same three heart function tests again.

Results showed that at rest, men and women showed little differences between the three tests. During the mental arithmetic task, all the volunteers showed an increase in heart rate and blood pressure, regardless of sex. However, while the men showed an increase in coronary vascular conductance under stress, the women showed no change.

## Taking Differences to Heart

This differing characteristic could potentially predispose women to heart problems while under stress, says study leader Chester Ray. He adds that the results came as a surprise, since previous studies men have significantly less blood flow than women during the physical stress of exercise, and could explain why women tend to have more heart troubles after stressful events, such as losing a spouse. The findings also reemphasize the importance of mental stress in affecting health.

"Stress reduction is important for anyone, regardless of gender," he explains, "but this study shines a light on how stress differently affects the hearts of women, potentially putting them at greater risk of a coronary event."

Further research, he says, could discern the mechanism behind this difference, leading to more targeted treatments and prevention efforts for [women](#) at risk of [coronary artery disease](#).

Provided by Federation of American Societies for

Experimental Biology

APA citation: Mental stress may be harder on women's hearts (2012, April 26) retrieved 5 May 2021 from <https://medicalxpress.com/news/2012-04-mental-stress-harder-women-hearts.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*