

Pre-op breast pain in about 28 percent of breast cancer patients

15 June 2012




Image courtesy of Blausen Medical

preoperative [breast pain](#)."

More information: [Abstract](#)

[Full Text \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

More than a quarter of women about to undergo breast cancer surgery experience breast pain, with genetic polymorphisms in inflammatory cytokines correlating with pain, according to a study published in the May issue of *The Journal of Pain*.

(HealthDay) -- More than a quarter of women about to undergo breast cancer surgery experience breast pain, with genetic polymorphisms in inflammatory cytokines correlating with pain, according to a study published in the May issue of *The Journal of Pain*.

Birha McCann, R.N., C.N.S., of the University of California in San Francisco, and associates recruited 398 women with breast cancer prior to surgery to determine the occurrence rate and characteristics of preoperative breast pain. The women completed self-report questionnaires and were genotyped. Variations in pro- and anti-inflammatory cytokine genes were assessed for those who did and did not report pain.

The researchers found that 28.2 percent of women self-reported breast pain. These women were significantly younger (P

"Findings suggest that breast pain occurs in over a quarter of women who are about to undergo [breast cancer surgery](#)," the authors write. "Based on phenotypic and genotypic characteristics found, inflammatory mechanisms contribute to

APA citation: Pre-op breast pain in about 28 percent of breast cancer patients (2012, June 15) retrieved 5 May 2021 from <https://medicalxpress.com/news/2012-06-pre-op-breast-pain-percent-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.