

PharmaNet system dramatically reduced inappropriate prescriptions of potentially addictive drugs

September 4 2012

A centralized prescription network providing real-time information to pharmacists in British Columbia, Canada, resulted in dramatic reductions in inappropriate prescriptions for opioid analgesics and benzodiazepines, widely used and potentially addictive drugs. The findings are reported in a study in *CMAJ (Canadian Medical Association Journal)*.

The study found that PharmaNet, a real-time prescription system implemented in BC pharmacies in July 1995, reduced potentially inappropriate prescriptions for opioids and benzodiazepines in two groups of patients—those on social assistance and seniors. The reductions ranged from about one-third (33%) to one-half (49%) depending on the drug and patient group.

"Our study demonstrates that a system like PharmaNet can help reduce the potentially inappropriate prescribing of medications that are prone to misuse," said lead author Colin Dormuth of the Therapeutics Initiative at the University of British Columbia. "In the five-year period of our study, the reduction in inappropriate refills was dramatic and sustained after the implementation of PharmaNet."

The PharmaNet system allows BC pharmacists to view the most recent 14 months of a patient's medication use regardless of which physician prescribed the drugs or which pharmacy dispensed them. The system

enables pharmacists to identify potentially harmful drug interactions, accidental duplications in therapy or potential prescription [drug misuse](#).

Researchers tracked "potentially inappropriate" prescriptions for opioids and benzodiazepines in BC, defined as a second prescription for the same drug issued by a different doctor and a different pharmacy within 7 days of a previous prescription for at least 30 tablets.

Benzodiazepines and [opioid analgesics](#) are [prescription medications](#) that are prone to misuse and addiction in some patients. Benzodiazepines include sedatives and anti-anxiety drugs such as diazepam, lorazepam and others, whereas [opioids](#) are taken for pain relief and include morphine, codeine, oxycodone and other drugs.

Most provinces have now adopted either a system allowing the [pharmacist](#) to view a patient's up-to-date medication history at the time of dispensing a medication or a real-time monitoring program specifically aimed at monitoring prescription drugs prone to misuse. Ontario adopted the latter approach in establishing its Narcotics Monitoring System earlier this year.

"Centralized drug information systems and monitoring programs are critical to help ensure these medications are used appropriately," observes coauthor David Juurlink, Head of Clinical Pharmacology and Toxicology at Toronto's Sunnybrook Health Sciences Centre and a scientist at the Institute for Clinical and Evaluative Sciences (ICES).

The study covered the period from Jan. 1, 1993 to Dec. 31, 1997—the 5-year period flanking the program's implementation. Because comprehensive data was available for people aged 65 years and over and patients receiving social assistance, the study focused on these two groups of patients.

More information: www.cmaj.ca/lookup/doi/10.1503/cmaj.120465

Provided by Canadian Medical Association Journal

Citation: PharmaNet system dramatically reduced inappropriate prescriptions of potentially addictive drugs (2012, September 4) retrieved 30 December 2022 from <https://medicalxpress.com/news/2012-09-pharmanet-inappropriate-prescriptions-potentially-addictive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.