

Patients use OTC NSAIDs even when they have a high risk of serious side effects

12 June 2013

A new study presented today at EULAR 2013, the Annual Congress of the European League Against Rheumatism, shows that one in eight patients at risk of developing a serious adverse drug event (ADE) is taking over-the-counter (OTC) non-steroidal anti-inflammatory drugs (NSAIDs), often to treat a musculoskeletal complaint.

Of these high risk OTC NSAID users, over one-third had taken the medication for more than 7 days, and 3% had exceeded the maximum recommended daily dosage.

Patients were considered at high risk of a serious ADE from OTC NSAIDs if they had a history of a [peptic ulcer](#) or ulcer complication, [myocardial infarction](#), stroke or heart failure, were aged over 70 years, had a [glomerular filtration rate](#)

Possible serious ADEs include [gastrointestinal bleeding](#), peptic ulceration, [high blood pressure](#) and worsening [heart failure](#).

"NSAIDs tend to be regarded by patients as harmless painkillers. However, in reality, even those available over the counter can cause a number of unpleasant side effects," said lead author of the study Aafke Koffeman of the Department Of General Practice, Erasmus Medical Center, Rotterdam, Netherlands.

"In most cases, it is likely to be ignorance of these potential ADEs rather than a deliberate disregard of the risks and contraindications", Ms Koffeman speculated. "These new data highlight the importance of healthcare professionals continuing to inform their patients of the risks of taking OTC NSAIDs, particularly where a new diagnosis or prescription increases their individual risk. High risk patients with painful musculoskeletal complaints should be advised to take safer alternative painkillers," she concluded.

Complications from the use of NSAIDs by high [risk](#)

[patients](#) are a common cause of unplanned [hospital admissions](#) related to medications, which in turn constitute a significant public health problem. Two previous Dutch observational studies have shown that approximately 5% of all unplanned hospital admissions are associated with ADEs, of which 40-46% are potentially preventable.²

This study was among adults registered with four Dutch general practitioners (GPs). Two samples of different patient populations were selected: (i) a random sample of adults (general population), and (ii) a sample of adult patients with a high risk of developing a serious ADE from NSAID use (high-risk population).

Patients' risk profiles were determined using their electronic medical records. All included patients were sent a questionnaire regarding their use of OTC NSAIDs in the four weeks prior to participation. In the high-risk population, of the 265 patients who agreed to participate, 33 (13%) had used an OTC NSAID. In the general population sample, of the 120 patients who agreed to participate, 35 (29%) had used an OTC NSAID.

Over 30% of OTC NSAID users had used the treatment for more than seven days in the high risk-population, and over 20% in the general population. OTC NSAIDs were used in a dosage exceeding the recommended daily maximum dose by 3% and 9% in the high-risk and the general populations respectively.

Musculoskeletal complaints formed the most commonly cited reason for OTC NSAID use in the high-risk group; 67% gave this as a reason for use.

More information: 1.Koffeman A et al., Use of over-the-counter non-steroidal anti-inflammatory drugs in the general population and in patients with a high risk of adverse drug events [abstract]. EULAR Annual European Congress of Rheumatology; 12-15 June 2013; Madrid, Spain.

Abstract nr. OP0202-PC.

2.Warlé-Van Herwaarden MF, Kramers C, Sturkenboom MC, et al. Targeting outpatient drug safety: Recommendations of the Dutch Harm-Wrestling Task Force. Drug Saf 2012; 5 (3): 245-59

Provided by European League Against Rheumatism

APA citation: Patients use OTC NSAIDs even when they have a high risk of serious side effects (2013, June 12) retrieved 5 May 2021 from <https://medicalxpress.com/news/2013-06-patients-otc-nsaids-high-side.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.