

Bariatric surgery restores nerve cell properties altered by diet

17 June 2013

Understanding how gastric bypass surgery changes the properties of nerve cells that help regulate the digestive system could lead to new treatments that produce the same results without surgery, according to Penn State College of Medicine scientists, who have shown how surgery restores some properties of nerve cells that tell people their stomachs are full.

The results may also better predict which patients will keep the weight off after surgery.

Roux-en-Y [gastric bypass](#) surgery is the most effective way to get severe obesity under control. Doctors make the stomach smaller and bypass a section of the small intestine. Besides restricting the amount of food a person can eat at one time, the procedure also seems to alter the properties of nerve cells.

"Restricting the size of the stomach has some role in the effectiveness of gastric bypass, but it's not the full story," said Kirsteen Browning, assistant professor of neural and behavioral sciences. "It is not fully understood why the surgery works." The researchers published their findings in the *Journal of Physiology*.

Complications from diseases such as [diabetes](#) can resolve before weight is lost, and sometimes before the person even leaves the hospital after gastric [bypass surgery](#).

"This suggests an altering of the [neural signals](#) from the gut to the brain and back," Browning said.

These nerve cells send signals to tell the body's [digestive system](#) how to respond properly and regulate normal functions of digestion. In obese people, the nerve cells are less excitable, meaning they respond less to normal stimulation. For example, there are [neurons](#) that help tell a person that their stomach is full, called satiation.

"These signals tell you to stop eating," said study co-author Andrew Hajnal, professor of neural and [behavioral sciences](#). "Obviously these signals are strong enough to be overcome by all of us and we can eat more even after we are told we are full. However, as [obesity](#) develops, it appears these signals are less strong and easier to overcome."

Penn State Hershey researchers used a high-fat diet in rats to replicate long-term exposure to a Western diet. They then observed the effects of gastric bypass on the rats and have shown for the first time that the effects of diet on nerve cells seem to be restored to normal function after the surgery. This would help in restoring satiation signals so that they can be recognized more easily.

"We know gastric bypass improves the health of nerve cells and reverses the effects on the signals," Browning said. "Even if the [nerve cells](#) have been affected over a long term, gastric bypass still improves toward normal function."

The goal of this line of research is to find new treatments that will not require gastric bypass surgery, which is still considered an invasive procedure.

"Once we understand what gastric bypass is doing, we hope we can mimic that with other treatments," Browning said. "Restricting the size of the [stomach](#) may not be the major player of the surgery; restoring normal neural function is also an important aspect."

Scientists may also be able to determine why the surgery is unsuccessful for some patients and better determine if surgery is the best option.

"Gastric bypass surgery doesn't work in all people," Hajnal said. "Some people regain the weight, but we don't know why. By understanding this happens, we may be able to predict in advance which patients will respond to the surgery."

Provided by Pennsylvania State University

APA citation: Bariatric surgery restores nerve cell properties altered by diet (2013, June 17) retrieved 6 May 2021 from <https://medicalxpress.com/news/2013-06-bariatric-surgery-nerve-cell-properties.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.