

Diagnosis of cervical spondylotic myelopathy delayed by primary care physicians

1 July 2013

Researchers from The Tel-Aviv Medical Center found delays in diagnosis of cervical spondylotic myelopathy (CSM) averaging 2 years in 42 patients who eventually underwent surgery. This is an important finding, because previous studies have shown that early diagnosis and treatment of CSM lead to better outcomes. The majority of patients initially sought a diagnosis for their symptoms from family physicians, who arrived at a correct diagnosis in only 4.8% of cases and never at the first clinic visit. Many other patients initially consulted community-based orthopedic surgeons, only one of whom suspected CSM at the first visit. Detailed findings of this study are reported and discussed in the article "Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians," by Eyal Behrbalk, M.D., Khalil Salame, M.D., Gilad J. Regev, M.D., Ory Keynan, M.D., Bronek Boszczyk, Dr. Med., and Zvi Lidar, M.D., published today, in the July 2013 issue of *Neurosurgical Focus*.

Cervical spondylotic myelopathy (CSM) has been called the most common form of spinal cord dysfunction in [older adults](#), although it can occur in younger people due to injury. As we age, vertebral joints in our necks begin to show wear (cervical spondylosis) and cervical spine ligaments thicken. These changes cause the [spinal canal](#) to narrow, which can lead to compression of the spinal cord. As spinal cord compression increases, [nerve cells](#) die, resulting in symptoms of cervical myelopathy that range from mild neck pain to motor weakness, [sensory loss](#), impaired walking, and, in cases of advanced disease, loss of bladder and [bowel control](#). CSM is a [progressive disease](#). Surgery is indicated when there is neurological impairment. Studies have shown that outcomes of surgery are best when the operation is performed during the early stages of the disease, when [neurological impairment](#) is minimal.

Between January 2009 and December 2010, 146 patients underwent surgery for CSM at The Spine Unit of Tel-Aviv Medical Center, a tertiary care center. All cases of degenerative pathological conditions of the cervical spine were included. Complete medical records (from both the Center and community-based physicians' offices) were available for 42 patients, and these patients comprised the study population. The researchers collected data from the time the first signs or symptoms of CSM were documented until the date of surgery. The diagnosis of CSM was based on the following: 1) symptoms consistent with CSM; 2) neurological findings suggestive of myelopathy; and 3) MRIs showing compression of the cervical spinal cord. In addition to the review of medical records, phone interviews with patients were conducted to obtain any missing clinical data.

Behrbalk and colleagues found that 69% of patients initially sought a diagnosis for their symptoms from family physicians and 21.4% from community-based orthopedic surgeons. The remaining patients consulted other physicians, none of whom was a neurologist or neurosurgeon, the most likely to recognize the disease. The researchers state that this follows the normal pattern of physician visits in Israel, where patients can directly schedule appointments with family physicians and orthopedic surgeons but require referrals to see neurologists or neurosurgeons. During the first physician visit, only one doctor—an orthopedic surgeon—suspected that CSM could be the underlying cause of the patient's symptoms.

At the second physician visits, most patients went to orthopedic surgeons (48.8%) and family physicians (26.8%); far fewer saw neurologists (9.8%) or neurosurgeons (2.4%). At the third physician visit, patients continued to consult orthopedic surgeons most frequently (38.5%), but

more patients sought out neurologists (25.6%) and neurosurgeons (18%) than before; 12.8% of patients returned to family physicians.

In this study, Behrbalk and colleagues found that it took a mean of 5.2 ± 3.6 physician visits to obtain the [correct diagnosis](#) of CSM. In the end, the diagnosis was made most often by neurosurgeons (38.1%) and neurologists (28.6%), and less frequently by orthopedic surgeons (19%), [family physicians](#) (4.8%), and a variety of other specialists (9.5%). The delays in diagnosis in this study, according to the researchers, rest primarily on a lack of awareness of CSM on the part of family practitioners and community-based orthopedic surgeons, who did not conduct full neurological examinations that could indicate myelopathy.

Given that CSM is a progressive disease, the researchers point out that delays in diagnosis can mean that "patients are referred to surgery at an advanced stage of the disease, at which point they are suffering from severe, often irreversible neurological damage."

Behrbalk and colleagues call for continued education of family practitioners and community-based [orthopedic surgeons](#) to make them more aware of CSM and promote early referrals for surgery when patients present with CSM symptoms.

More information: Behrbalk E, Salame K, Regev GJ, Keynan O, Boszczyk B, Lidar Z: Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurgical Focus* 35 (1):E1, July 2013; [DOI: 10.3171/2013.3.FOCUS1374](#)

Provided by Journal of Neurosurgery

APA citation: Diagnosis of cervical spondylotic myelopathy delayed by primary care physicians (2013, July 1) retrieved 11 June 2022 from <https://medicalxpress.com/news/2013-07-diagnosis-cervical-spondylotic-myelopathy-primary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.