

More awareness, fast response key to combatting stroke in children

12 February 2014

Parents and healthcare professionals must be aware that children can have strokes and be prepared to respond to symptoms, according to research presented at the American Stroke Association's International Stroke Conference 2014.

As in adults, warning signs of [stroke](#) in children are: sudden weakness or numbness of the face, arm or leg; sudden difficulty in speaking; sudden problems in seeing; sudden difficulty walking; dizziness; or sudden onset of headache.

Researchers interviewed 28 parents whose child had a stroke about factors that contributed to delayed arrival to the hospital. The children were infants to teens. Symptoms of stroke didn't vary by age, although seizures were more common in younger children and strokes mostly occurred at home and to a lesser extent at school.

Though most of the parents thought the symptoms were serious, only about half called 9-1-1; 36 percent considered the possibility of stroke; and 21 percent had a "wait and see" attitude or called a relative before taking emergency action.

"Think stroke, act fast, and call 9-1-1," said Mark MacKay, M.D., lead researcher and director of the Children's Stroke Program at the Royal Children's Hospital and Murdoch Children's Research Institute in Melbourne, Australia. "That message applies to adults and children."

Median time from [symptom onset](#) to emergency room arrival was 1.8 hours, with some taking as long as 24 hours.

In previous studies in the United States, Canada, England and Australia, the average time delay to diagnose stroke in children was more than 24 hours.

"Getting to the hospital quickly is an essential first

step to develop strategies to improve access to emergency treatment in children," MacKay said.

Children won't receive the benefits of emergency clot-busting treatment unless "we develop strategies to improve the rapid recognition of [childhood stroke](#)," he said. Clot-busting treatment must be started within three to four and a half hours of symptom onset.

Child neurologists should educate primary care physicians, pediatricians, heart specialists and emergency room physicians about stroke in children and parents of children with conditions associated with increased risk such as [sickle cell disease](#) or heart conditions urgently need to be aware, MacKay said.

In the United States, 4.6 per 100,000 children (0 to 19 years) had a stroke in 1997-2003, according to the American Heart Association. Despite treatment, one of 10 [children](#) with ischemic or [hemorrhagic stroke](#) will have a recurrence within five years.

Provided by American Heart Association

APA citation: More awareness, fast response key to combatting stroke in children (2014, February 12) retrieved 17 October 2022 from <https://medicalxpress.com/news/2014-02-awareness-fast-response-key-combatting.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.