

Anesthetic technique improves quality of recovery for women having breast cancer surgery

March 11 2014

Anesthesiologists using a technique similar to a dental freeze can improve the quality of recovery and decrease recovery time for breast cancer surgery patients, according to a new study.

The study, from researchers at St. Michael's Hospital and Women's College Hospital, was published in the March edition of *Anesthesiology*. It is the world's first randomized control trial for [breast cancer](#) surgery that compares the use of ultrasound-guided paravertebral blocks – a [local anesthetic](#) freezing that blocks breast nerves – to general anesthetic. The findings reveal that [breast cancer patients](#) who received local anesthetic had superior pain relief, spent less time in recovery rooms after surgery, and were discharged an hour earlier than patients who were put under [general anesthesia](#).

"Real time, image-guided ultrasound nerve blocks have revolutionized the practice of regional anesthesia," said Dr. Faraj Abdallah, an anesthesiologist at St. Michael's Hospital and lead author of the study. "This is the first study to show how effective these ultrasound-guided blocks can now be for breast [cancer surgery](#). Even more importantly, we've been able to demonstrate that blocks help patients feel better and return to their normal levels of mental and physical functionality sooner after surgery."

Dr. Abdallah conducted this randomized controlled trial of 64 women at

Women's College Hospital, the first Canadian site to complete a study where this image-guided technique was used in breast cancer surgery. Because of its demonstrated benefits and the addition of Dr. Abdallah, St. Michael's Hospital is working on incorporating this technique into practice in its CIBC Breast Centre.

The paravertebral block technique uses ultrasound to precisely guide a needle to intercostal nerves reaching the breast and deliver local anesthetic to freeze these nerves. In total, five blocks are needed to freeze the five nerves involved. The freeze blocks the transmission of pain from the breast where surgeons can perform full or partial mastectomies, or even reconstructive breast surgery. At the same time, patients are given an intravenous drug called propofol which allows them to sleep through the surgery and continue breathing on their own, without needing a breathing tube or a ventilator.

"There is a global interest in avoiding general anesthesia in breast cancer surgery whenever possible because of potential long-term effects. Although effective, general anesthesia use is also associated with nausea, vomiting and can result in excessive sedation even after surgery," said Dr. Tulin Cil, a study co-author and general surgeon at Women's College Hospital. "Patients who receive general anesthesia also require extra help to look after themselves in the first few days after surgery, whereas patients who received the nerve blocks are much more independent and require less support."

In the past, paravertebral blocks were inserted "blindly" – without the use of guided ultrasound – and, not surprisingly, with a high failure rate.

"Without ultrasound it may take several attempts to administer the blocks," said Dr. Abdallah. "And even then, it would be difficult to tell if they'd hit their mark so [patients](#) might feel pain during surgery and would then need to be put under general anesthetic, which defeats the

purpose."

Dr. Abdallah said that the benefits of paravertebral blocks following [breast cancer surgery](#) would also apply to cosmetic breast surgeries such as reductions or augmentation because the same breast nerves can be frozen in a similar manner.

Provided by St. Michael's Hospital

Citation: Anesthetic technique improves quality of recovery for women having breast cancer surgery (2014, March 11) retrieved 29 February 2024 from <https://medicalxpress.com/news/2014-03-anesthetic-technique-quality-recovery-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.