

Statins may reduce risk of progression of renal cancer

April 23 2014



(HealthDay)—Use of statins is associated with a reduced risk of progression of localized renal cell carcinoma, according to research published in the April issue of *The Journal of Urology*.

In an effort to assess the effect of statin medications on the progression of the neoplasm, Robert J. Hamilton, M.D., M.P.H., of the Memorial Sloan-Kettering Cancer Center in New York City, and colleagues analyzed data for 2,608 patients receiving surgical treatment for localized renal cell carcinoma.

The researchers found that 27 percent of the patients undergoing surgery for localized renal cell carcinoma were statin users. At a median follow-

up of 36 months, statin use was associated with a significant reduction in the risk of progression of localized renal cell carcinoma following surgery (hazard ratio [HR], 0.67; 95 percent confidence interval [CI], 0.47 to 0.96; $P = 0.028$) and a nonsignificant reduction in overall mortality (HR, 0.89; 95 percent CI, 0.71 to 1.13; $P = 0.3$). When statin use was modeled as a time-dependent covariate, the reduction in risk of progression of disease was decreased (HR, 0.77; $P = 0.12$) and the reduction in risk of overall mortality was increased (HR, 0.71; $P = 0.002$).

"This is the first study to directly examine the association between [statin](#) use at surgery and progression to metastasis or [renal cell carcinoma](#) death," the authors write.

More information: [Abstract](#)
[Full Text](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Statins may reduce risk of progression of renal cancer (2014, April 23) retrieved 7 April 2023 from <https://medicalxpress.com/news/2014-04-statins-renal-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--