

Sleep apnea tied to hearing loss in large study

20 May 2014




Illustration of obstruction of ventilation. Credit: Habib M'henni / public domain

Both high and low frequency hearing impairment have been linked with sleep apnea in a new study of nearly 14,000 individuals.

"In our population-based study of 13,967 subjects from the Hispanic Community Health Study/Study of Latinos, we found that sleep apnea was independently associated with hearing impairment at both high and low frequencies after adjustment for other possible causes of [hearing loss](#)," said lead author Amit Chopra, MD, currently at the Albany Medical Center in New York.

The study was presented at the 2014 American Thoracic Society International Conference.

All subjects underwent successful in-home sleep apnea studies and on-site audiometric testing at

baseline. Sleep apnea was assessed with the apnea-hypopnea index (AHI), which indicates sleep apnea severity based on the number of apnea (complete cessation of airflow) and hypopneas (partial cessation of airflow) per hour of sleep. Sleep apnea was defined as an AHI \geq 15 events/hour. High frequency hearing impairment was defined as having a mean hearing threshold of greater than 25 decibels in either ear at 2000, 3000, 4000, 6000 and 8000 Hz, and low frequency hearing impairment was defined as having a mean hearing threshold of greater than 25 decibels in either ear at 500 Hz and 1000 Hz.

Among the 13,967 study subjects, 9.9% had at least moderate sleep apnea (AHI \geq 15), 19.0% had high frequency hearing impairment, 1.5% had low frequency hearing impairment, and 8.4% had both high and low frequency hearing impairment.

Hearing impairment was more common among individuals of Cuban and Puerto Rican descent and among those with a higher body mass index, self-reported snoring and/or sleep apnea. Sleep apnea was independently associated with a 31% increase in high frequency hearing impairment, a 90% increase in low frequency hearing impairment, and a 38% increase in combined high and low frequency hearing impairment in analyses adjusting for age, sex, background, history of hearing impairment, external noise exposure, conductive hearing loss and other factors. Increasing AHI was associated with a higher prevalence of [high frequency](#), but not [low frequency](#) hearing impairment.

A limitation of this study was that the authors could not account for the impact of treatment of sleep apnea on their findings. However, it was rare for the participants to be on sleep apnea treatment in this cohort.

"Patients with sleep apnea are at increased risk for a number of comorbidities, including heart disease

and diabetes, and our findings indicate that sleep apnea is also associated with an increased risk of hearing impairment" said Dr. Chopra. "The mechanisms underlying this relationship merit further exploration. Potential pathways linking sleep apnea and [hearing impairment](#) may include adverse effects of [sleep apnea](#) on vascular supply to the cochlea via inflammation and vascular remodeling or noise trauma from snoring ."

Provided by American Thoracic Society

APA citation: Sleep apnea tied to hearing loss in large study (2014, May 20) retrieved 4 December 2022 from <https://medicalxpress.com/news/2014-05-apnea-tied-loss-large.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.