

Low cholesterol linked with worse survival in patients with kidney cancer

12 June 2014

People are often told to reduce their cholesterol to improve their heart health, but new research suggests that low cholesterol may increase kidney cancer patients' risk of dying from their disease. The findings, which are published in *BJU International*, indicate that cholesterol testing may help doctors as they monitor and treat patients with kidney cancer.

patients with renal cell carcinoma." Michela de Martino, Carmen V. Leitner, Christoph Seemann, Sebastian L. Hofbauer, Ilaria Lucca, Andrea Haitel, Shahrokh F. Shariat, and Tobias Klatte. *BJU International*; Published Online: June 12, 2014 (DOI: 10.1111/bju.12767).

Increasing evidence suggests that alterations in cholesterol and other lipids are associated with the development, progression, and prognosis of various cancers. To assess the situation as it relates to kidney cancer, Tobias Klatte, MD, of the Medical University of Vienna in Austria, and his colleagues analyzed total blood cholesterol levels in 867 patients with renal cell carcinoma before they underwent kidney surgery. The investigators then followed the patients for a median of 52 months.

Provided by Wiley

Low <u>blood cholesterol</u> before treatment was associated with more advanced tumor stages and cancer spread during follow-up. Also, patients with high cholesterol had a 43 percent lower risk of dying from their cancer compared with patients with low cholesterol. Finally, including patients' cholesterol levels with traditional risk factors increased the accuracy of prognoses.

It's unclear how cholesterol may affect a kidney cancer patients' prognosis. It may be that certain components of cholesterol impact the activity of cancer-related pathways to affect tumor growth and spread.

"As this was a hypothesis-generating study, our findings should be confirmed in independent datasets. If confirmed, patients with <u>low cholesterol</u> may be considered high-risk and may be treated or followed up more aggressively," said Dr. Klatte.

More information: "Preoperative serum cholesterol is an independent prognostic factor for



APA citation: Low cholesterol linked with worse survival in patients with kidney cancer (2014, June 12) retrieved 4 September 2022 from https://medicalxpress.com/news/2014-06-cholesterol-linked-worse-survival-patients.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.