

No correlation between baby formulas and development of diabetes-associated autoantibodies

June 16 2014

There is no correlation between the consumption of a cow's milk-based formula or hydrolyzed protein formula and the development of diabetes-associated autoantibodies in children younger than seven, according to a worldwide research study published in the *Journal of the American Medical Association (JAMA)*.

The Trial to Reduce Insulin Dependent Diabetes Mellitus in the Genetically at Risk (TRIGR) study, which examines possible reduction in autoantibodies associated with type 1 diabetes, tracked children genetically predisposed to type 1 diabetes for seven years.

It was conducted in 78 centers in 15 countries, including the U.S and Canada, and will continue to receive funding from the National Institutes of Health to collect data up to the children's 10th birthday. A pilot study in Finland with a smaller, homogeneous population showed that weaning infants to an extensively hydrolyzed casein formula decreased the cumulative occurrence of diabetes-associated autoantibodies in children who have a first-degree relative with type 1 diabetes and an increased genetic risk.

Anita Nucci, assistant professor of nutrition at Georgia State University, is the U.S. coordinator and the North America nutrition coordinator for TRIGR. She notes that while there is no correlation between cow's milk protein and the [autoantibodies](#) that result in type 1 diabetes, the second

end-point of the TRIGR study is the development of type 1 diabetes by 10 years of age.

"The NIH's continued support validates the possibility the connection with cow's [milk protein](#) and type 1 diabetes may appear a little later in the lives of children who live outside Finland or that hydrolyzed formula affects the rate of progression of autoimmunity to [clinical diabetes](#) in high-risk children," says Nucci.

Some genetic subgroups may carry a higher risk for diabetes and cow's milk proteins. Scandinavia is the region with the highest incidence of [type 1 diabetes](#) in the world.

Provided by Georgia State University

Citation: No correlation between baby formulas and development of diabetes-associated autoantibodies (2014, June 16) retrieved 15 July 2023 from <https://medicalxpress.com/news/2014-06-baby-formulas-diabetes-associated-autoantibodies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.