

TASTE trial finds no benefit of thrombus aspiration for AMI at one year

2 September 2014

Thrombus aspiration for acute myocardial infarction (AMI) does not reduce mortality or other clinical endpoints long term, according to the awaited one-year follow-up results from the Thrombus Aspiration in ST- Elevation myocardial infarction in Scandinavia (TASTE) trial. The findings were presented at ESC Congress today by Dr Bo Lagerqvist from Sweden.

Dr Lagerqvist said: "The results of the one-year follow-up of the TASTE trial challenge the current practice of blood clot aspiration after a <u>heart attack</u> ."

TASTE was a multicentre, prospective, randomised, controlled open-label trial that enrolled 7 244 patients from Sweden, Denmark and Iceland who had a diagnosis of ST-elevation <u>myocardial infarction</u> (STEMI). Half of the patients were assigned to balloon treatment only (known as <u>percutaneous coronary intervention</u>, or PCI) and the other half had their blood clot aspirated before PCI.

As previously reported from the TASTE trial, the death rate when the patients were followed 30 days after treatment was not lower after clot aspiration. Today the investigators report the results after patients have been followed for one year.

The results presented today are consistent with the previous report. After one year the death rate was not statistically different between the two treatment groups (5.3% after clot aspiration and 5.6% after PCI alone). Similarly, there was no difference between the two groups regarding other endpoints including the risk of new heart attack, stroke and complications related to the treatment. Even high risk groups such as smokers, patients with diabetes or patients with large clots had similar results with either approach.

Dr Lagerqvist said: "Our findings do not support a

role for this clot aspiration as a routine additional treatment in heart attacks."

He added: "Current ESC guidelines on treatment of patients with STEMI recommend that thrombus aspiration should be considered and most opinion leaders advocate its use. Since the TAPAS trial suggested a mortality benefit after one year thrombus aspiration has gained enormous popularity. The therapy is so popular among interventional cardiologists because it intuitively feels beneficial to aspirate the clot that closes the artery."

Dr Lagerqvist continued: "The TASTE trial is the first large-scale <u>randomised trial</u> of thrombus aspiration for STEMI to be large enough to reveal meaningful findings on mortality and morbidity. It enrolled more patients than all previous randomised <u>trials</u> of this procedure combined and included a much broader range of patients in order to make the results relevant to everyday clinical practice."

The study's unique Registry-Based Randomised Clinical Trial (RRCT) protocol used national registries as on-line platforms for randomisation, case record forms and follow-up, making the trial economically and administratively feasible.

Study chair Professor Stefan James (Sweden) said: "TASTE is the first trial ever to use the registry based trial concept. This concept enabled enrolment of more than 80% of all eligible patients nationwide with complete control of the nonenrolled <u>patients</u> and unlimited duration of follow-up at a very low cost."

More information: "ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation." *European Heart Journal.* 2012;33:2569–2619. DOI: 10.1093/eurheartj/ehs215



Svilaas T, Vlaar PJ, van der Horst IC, Diercks GF, de Smet BJ, van den Heuvel AF, Anthonio RL, Jessurun GA, Tan ES, Suurmeijer AJ, Zijlstra F. "Thrombus aspiration during primary percutaneous coronary intervention." *N Engl J Med*. 2008;358(6):557-567. DOI: 10.1056/NEJMoa0706416.

Provided by European Society of Cardiology

APA citation: TASTE trial finds no benefit of thrombus aspiration for AMI at one year (2014, September 2) retrieved 19 September 2022 from <u>https://medicalxpress.com/news/2014-09-trial-benefit-thrombus-aspiration-ami.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.