

Spinal cord stimulation feasible for diabetic neuropathy

30 September 2014



Image courtesy of Blausen Medical

The study was funded by Medtronic.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

(HealthDay)—For patients with painful diabetic peripheral neuropathy (PDPN), spinal cord stimulation (SCS) is a successful treatment, according to a study published online Sept. 11 in *Diabetes Care*.

Rachel Slangen, from the Maastricht University Medical Centre in the Netherlands, and colleagues randomized 36 PDPN patients to SCS in combination with best medical treatment (BMT; 22 patients) and BMT only (14 patients). If trial stimulation was successful, the SCS system was implanted. Definitions of [treatment success](#) include ≥50 percent pain relief during daytime or nighttime or "(very) much improved" for pain and sleep on the patient global impression of change scale at six months.

The researchers found that 59 percent of the SCS and 7 percent of BMT met the definition of treatment success (P pain relief were reported by 41 and 36 percent in the SCS group, respectively, and by 0 and 7 percent in the BMT group, respectively (P

"Treatment success was shown in 59 percent of [patients](#) with PDPN who were treated with SCS over a six-month period, although this treatment is not without risks," the authors write.

APA citation: Spinal cord stimulation feasible for diabetic neuropathy (2014, September 30) retrieved 26 September 2022 from <https://medicalxpress.com/news/2014-09-spinal-cord-feasible-diabetic-neuropathy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.