

# Kidney failure later in life can be predicted based on measurements taken in adolescence

5 November 2014

Researchers at Örebro University found that high blood pressure, being overweight, elevation of a blood marker indicating inflammation, or the presence of protein in the urine in otherwise healthy teenage males were all independent predictors of end stage renal disease (ESRD) in later in life.

"We were surprised by the magnitude of the risk, especially for individuals with unexplained proteinuria (protein in urine) in adolescence," said Dr. Per-Ola Sundin, lead author of the study and a researcher and physician at Örebro University Hospital. "This shows that silent kidney damage is already occurring in teenagers who won't develop more noticeable symptoms of kidney disease for many years."

The study was based on Swedish male residents who had a physical examination at the time they were compulsorily enlisted in the military in the early 1970s; when the majority of these individuals were 18 and 19 years old. Up to the age of 58 years, researchers identified 534 men who developed ESRD and compared them to a similar group of 5127 men who did not develop ESRD.

The presence proteinuria in adolescence was associated with a seven-fold increase in the risk of ESRD. Low-grade inflammation found by testing for the [erythrocyte sedimentation rate](#) in blood indicated a two-fold increase in kidney failure risk. Hypertension, defined as [blood pressure](#) of 160/100 and higher, quadrupled the risk of developing ESRD, while a Body Mass Index (BMI) of 30 or above (indicating obesity) increased the ESRD risk by 3.5 times. Although the study was limited to Swedish men, researchers stated there was no biological reason that these findings would not be relevant to females and populations in other industrialized countries.

This study highlights the importance of adopting a life-course approach to understanding how risks for serious kidney disease begin to accumulate many years before the disease is diagnosed. "These results suggest that there is a screening opportunity at an early age if we want to prevent chronic [kidney disease](#)," said Dr. Sundin. "Particularly, proteinuria without an explanation should be followed up by doctors and patients. Also, it is never too early to tackle issues like unhealthy weight gain and hypertension as it may be much harder to begin a healthy lifestyle in later adulthood. How you live as a teenager has consequences for your health in middle age".

**More information:** "Predictors in Adolescence of ESRD in Middle-Aged Men." *American Journal of Kidney Diseases*, April 1, 2014; Accepted: June 9, 2014; Published Online: August 12, 2014. [dx.doi.org/10.1053/j.ajkd.2014.06.019](http://dx.doi.org/10.1053/j.ajkd.2014.06.019)

Provided by Örebro Universitet

APA citation: Kidney failure later in life can be predicted based on measurements taken in adolescence (2014, November 5) retrieved 4 June 2021 from <https://medicalxpress.com/news/2014-11-kidney-failure-life-based-adolescence.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*