


n-3 fatty acids cut nonesterified fatty acid, T2DM link

20 January 2015



(HealthDay)—Nonesterified fatty acids (NEFAs) are associated with type 2 diabetes, and the association is modified by n-3 FA levels, according to a study published online Jan. 8 in *Diabetes Care*.

Brian T. Steffen, Ph.D., from the University of Minnesota in Minneapolis, and colleagues examined whether [serum levels](#) of NEFAs relate to the risk of incident [type 2 diabetes](#), and whether plasma n-3 FAs modify this correlation. The authors measured NEFAs in fasting serum in 5,697 participants of the Multi-Ethnic Study of Atherosclerosis, and determined phospholipid n-3 FAs eicosapentaenoic and docosahexaenoic acids in plasma.

The researchers identified higher diabetes incidence across successive NEFA quartiles over a mean 11.4-year study period ($P_{\text{trend interaction}} = 0.03$). A higher risk of type 2 diabetes was seen across quartiles of NEFAs ($P_{\text{trend}} < 0.001$) among individuals with lower n-3 levels (

APA citation: n-3 fatty acids cut nonesterified fatty acid, T2DM link (2015, January 20) retrieved 11 October 2022 from <https://medicalxpress.com/news/2015-01-n-fatty-acids-nonesterified-acid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no

part may be reproduced without the written permission. The content is provided for information purposes only.