

A new perspective on Phantom Eye Syndrome

29 May 2015



Researchers from the University of Liverpool have found that approximately half of patients who have an eye removed because of a form of eye cancer experience 'phantom eye syndrome.'

Patients with the condition experience "seeing" and pain in the eye that is no longer there. Researchers assessed 179 [patients](#) whose eye had been removed as a result of a cancer, called intraocular melanoma.

They found that more than a third of the patients experienced phantom eye symptoms every day. In most patients, the symptoms ceased spontaneously, but some patients reported that they have to do something to stop the sensation, such as distracting themselves or blinking.

Pattern and colours

The 'phantom' sensations often start several weeks after the eye is removed, eventually subsiding over time. Most see only patterns or colours, but some feel they see people and scenes.

More than one-in-four reported that they sometimes feel that they can even see what is actually happening around them. A similar number feel pain in the non-existent eye.

The researchers found that a fifth of patients find these sensations pleasurable, but a similar number are disturbed by them.

Largest study

Health psychologist Laura Hope-Stone, from the University's Institute of Psychology, Health and Society (IPHS) who led the research, said: "To our knowledge, this is the largest study of phantom symptoms in patients who have lost an eye for intraocular melanoma.

"We carried out the study because many patients have told me about symptoms like this when I see them after surgery for psychological support, but we did not know exactly how common they are."

Health psychologist, Dr Steve Brown, also from IPHS, added: "The size of the study means we can now tell whether certain kinds of people are more likely to have phantom symptoms.

"They are most common in younger patients, and having pain in the non-existent eye is more likely in patients who are anxious or depressed, although we don't yet know why that's the case."

No guarantee

Professor Heinrich Heimann, who leads the Liverpool Ocular Oncology Centre at the Royal Liverpool University Hospital where the study was conducted, and the largest of the four UK centres which treat ocular melanoma, said: "These results are important because now we can tell patients not to be alarmed if they experience visual sensations or pain afterwards.

"We can also warn them that, unfortunately, removing an [eye](#) does not guarantee that there will be no [pain](#) weeks or months afterwards."

More information: "Phantom Eye Syndrome."
[dx.doi.org/10.1016/j.optha.2015.04.005](https://doi.org/10.1016/j.optha.2015.04.005)

Provided by University of Liverpool

APA citation: A new perspective on Phantom Eye Syndrome (2015, May 29) retrieved 11 June 2021 from <https://medicalxpress.com/news/2015-05-perspective-phantom-eye-syndrome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.