

To what extent are condoms responsible for erection difficulties?

18 August 2015



Image: Wikipedia

among Young, Heterosexual Men Who Do and Do Not Report Condom-Associated Erection Problems (CAEP)." *The Journal of Sexual Medicine* [DOI: 10.1111/jsm.12964](https://doi.org/10.1111/jsm.12964)

Provided by Wiley

Research indicates that the use of condoms may cause some men to experience erection difficulties. However, in a study of 479 heterosexual men who used condoms and were 18 to 24 years old, those who reported condom-associated erection problems were also more likely to experience more generalized erection difficulties. Investigators also found that more than one-third of participants had never been taught how to use a condom correctly.

Clinicians should assess whether men using condoms experience condom-associated erection problems and where appropriate, make referrals for psychosexual therapy or provide educational information on condom use.

"Condom-associated erection problems have been a very under-researched topic. Increasing evidence suggests, however, that they may influence whether [condoms](#) are used correctly or from start to finish of sex," said Dr. Cynthia Graham, co-author of *The Journal of Sexual Medicine* study.

More information: "General Erectile Functioning

APA citation: To what extent are condoms responsible for erection difficulties? (2015, August 18)
retrieved 2 August 2022 from <https://medicalxpress.com/news/2015-08-extent-condoms-responsible-erection-difficulties.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.