

# Antiarrhythmia drugs no impact on late AF recurrence

30 August 2015

In patients with atrial fibrillation (AF) treated with radiofrequency catheter ablation, the addition of antiarrhythmic drugs (AADs) for 90 days after the procedure did not reduce arrhythmia recurrence rates at one year, according to results of the Efficacy of Antiarrhythmic Drugs Short-Term Use after Catheter Ablation for Atrial Fibrillation (EAST-AF) trial.

Results of the study, presented as a Hot Line at ESC Congress 2015.

The study did show a temporary benefit of AAD, but the effect disappeared as soon as medication was stopped, suggesting no benefit to making post-ablation AAD therapy standard practice, said study investigator Kazuaki Kaitani, MD, from Tenri Hospital in Tenri, Nara, Japan.

Radiofrequency ablation around the pulmonary veins, known as "pulmonary vein isolation" (PVI), is the standard treatment for AF, but recurrence is common.

Short-term (6 weeks) therapy with AADs after PVI for paroxysmal AF has been previously shown to reduce early arrhythmia recurrence rates compared to no AADs, but this benefit disappeared after treatment discontinuation

The EAST-AF trial investigated whether a longer treatment period might extend the benefit beyond AAD discontinuation. The study included 2,044 patients (mean age 63 years) undergoing first radiofrequency catheter ablation for AF at 19 cardiovascular centres in Japan.

Patients were randomised to treatment with AADs (n=1,016) or a [control group](#) (n=1,022) for 90 days after the procedure.

The primary endpoint of the study was arrhythmias lasting for more than 30 seconds or requiring repeat ablation, hospital admission, cardioversion

or AADs between 3 months and 1-year post ablation.

While there was a significantly lower rate of early arrhythmia recurrences in the AAD group compared with the control group during the treatment period (41.0% versus 47.9%, respectively; P=0.013), this benefit did not persist after AAD discontinuation.

The 1-year recurrence-free rate was 69.5% in the AAD group and 67.8% in the control group, with no significant difference between the two groups (adjusted hazard-ratio [HR] 0.93; 95% confidence-interval, 0.79 to 1.09; P=0.38).

In addition, this finding was consistent across all prespecified subgroups, including patients with paroxysmal AF versus those with persistent or long-lasting AF.

"Therefore, we concluded that the short-term use of AADs for 90-days following [catheter ablation](#) of AF reduces the incidence of recurrent AF during the 90-day treatment period, but does not lead to improved clinical outcomes in the later phase," he said.

Provided by European Society of Cardiology

APA citation: Antiarrhythmia drugs no impact on late AF recurrence (2015, August 30) retrieved 6 May 2021 from <https://medicalxpress.com/news/2015-08-antiarrhythmia-drugs-impact-late-af.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*