


Signaling pathway suppresses brain tumors

4 December 2015



Cross-section of a murine stem cell-derived brain tumor with histological features strikingly similar to human gliomas. Credit: University of Basel, Claudio Giachino/Verdon Taylor

Researchers at the University of Basel took a close look at a signaling pathway present in most organisms and found that it suppresses the formation of specific types of brain tumor. Their results have been published by the journal *Cancer Cell*.

Gliomas are the most common [brain tumors](#) in adults and the prognosis for patients is, in many cases, very bad. Therefore, novel and effective therapies for glioma treatment are needed. In order to develop these, it is crucial to understand the biology of this type of tumor.

Stem cells as potential source of tumors

So far it has been highly debated which brain cells can form gliomas when they acquire gene mutations. However, researchers believe that [brain stem cells](#) might be a potential source of this type of cancer. Stem cells in the human brain can generate new nerve cells and, if something goes wrong in this process and uncontrolled proliferation or impaired differentiation occurs, this may lead to the formation of a brain tumor.

A research team led by Professor Verdon Taylor from the Department of Biomedicine at the

University of Basel has now studied whether one molecular mechanism that controls normal stem cell maintenance in the brain is hijacked and used by cancer [cells](#) during [tumor formation](#).

Active signaling pathway suppresses tumor formation

The researchers studied the so-called Notch pathway. This [signaling pathway](#) is central to [brain stem cell](#) activity and it has been proposed to – once aberrantly activated – contribute to the growth of gliomas. "In contrast to our expectations, we found that the opposite is the case: when activated, this pathway actually suppresses the formation of some types of glioma", says Claudio Giachino, first author of the study. Conversely, in some forms of glioma the inactivation of the pathway results in accelerated growth and makes the tumor more aggressive.

Due to these properties, the Notch pathway could, in the future, not only serve as a new therapeutic target but could also be used as a new diagnostic tool in order to get more reliable prognoses for disease progression and patient survival. "Our results demonstrate major differences in the molecular requirements between seemingly similar types of [brain](#) tumor and indicate that gliomas must be carefully examined before selecting potentially specific therapeutic interventions in the future", says Taylor.

More information: Claudio Giachino, et al. A Tumor Suppressor Function for Notch Signaling in Forebrain Tumor Subtypes. *Cancer Cell* (2015), DOI: [10.1016/j.ccell.2015.10.008](https://doi.org/10.1016/j.ccell.2015.10.008)

Provided by University of Basel

APA citation: Signaling pathway suppresses brain tumors (2015, December 4) retrieved 1 May 2021 from <https://medicalxpress.com/news/2015-12-pathway-suppresses-brain-tumors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.