

Treating major depression in older adults with diabetes may lower risk of death

27 January 2016

According to a new study published in the *Journal* of the American Geriatrics Society, effective treatment for depression could go a long way toward improving health status and even preventing death among older adults who also have diabetes.

Researchers assessing depression care managers working with physicians and older adults as reducing the suffering produced by depression. to offer recommendations and ensure followthrough on those suggestions—found that older people with diabetes who received structured depression care management were 53 percent less likely to die over the course of the study when compared with people who received care as usual. The researchers concluded that depression care management also seemed to reduce the risk of death for people with other chronic conditions (except heart disease), but not to the same extent as what they saw with diabetes.

Depression has a number of effects on health, including changing how the body maintains balance and reacts to stress and making it harder for people to be physically active, engage in selfcare, and follow through on treatment plans. While these risks have been known for some time, it has been less clear whether treating depression would translate into improvements in health.

In this new study, researchers evaluated 1,226 people over the age of 60 who had been diagnosed with major depression, minor depression, or no depression. Participants completed questionnaires about their health, including whether or not they had chronic diseases such as diabetes or heart disease.

In addition to the reduction in mortality, the study also found that the structured treatment improved depression considerably more than did care as usual. Forty percent of the participants experienced complete relief of depression

symptoms after four months of treatment, compared to 22.5 percent of older people receiving care as usual.

As one of the first studies to examine if treating depression reduces the risk for death across chronic medical conditions, this work highlights the need to integrate mental health and primary care management—consisting of trained depression careservices. Doing so may end up saving lives as well

> More information: Hillary R. Bogner et al. Does a Depression Management Program Decrease Mortality in Older Adults with Specific Medical Conditions in Primary Care? An Exploratory Analysis, Journal of the American Geriatrics Society (2016). DOI: 10.1111/jgs.13711

Provided by American Geriatrics Society

1/2



APA citation: Treating major depression in older adults with diabetes may lower risk of death (2016, January 27) retrieved 28 August 2022 from https://medicalxpress.com/news/2016-01-major-depression-older-adults-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.